

Open-closed field algebras

Liang Kong

Abstract

We introduce the notions of open-closed field algebra and open-closed field algebra over a vertex operator algebra V . In the case that V satisfies certain finiteness and reductivity conditions, we show that an open-closed field algebra over V canonically gives an algebra over a $\mathbb{C}$ -extension of Swiss-cheese partial operad. We also give a tensor-categorical formulation and constructions of open-closed field algebras over V .

0 Introduction

Open-closed (or boundary) conformal field theory was first developed in physics by Cardy [C1]-[C4][CL]. It has important applications to certain critical phenomena on surfaces with boundaries in condensed matter physics. It is also a powerful tool in the study of “D-branes” in string theory. A mathematical definition of such theory, which generalizes Segal’s definition of conformal field theory [S1][S2], is given by Huang [H7], Hu and Kriz [HKr]. However, a thorough mathematical study of this subject is still lacking.

In [V], Voronov introduced the so-called “Swiss-cheese operad” to describe the interactions among open strings and closed strings. In order to incorporate the full conformal symmetry which is not included in Voronov’s notion of Swiss-cheese operad, Huang and the author introduced in [HKo1] a notion called Swiss-cheese partial operad. One of the main goals of this work is to construct examples of algebras over certain $\mathbb{C}$ -extension of Swiss-cheese partial operad.

Swiss-cheese partial operad consists of disks with strips and tubes, which are conformal equivalence classes of disks with oriented punctures in the interior of the disks (called tubes) and those on the boundaries of the disks (called strips), together with a local coordinate map around each puncture. On the moduli space of disks with strips and tubes, which is denoted as $\mathfrak{S}$, there are two kinds of sewing operations. One is called boundary sewing operation, which sews two elements in $\mathfrak{S}$ along two oppositely oriented strips. The other one is called interior sewing operation, which sews a disk with strips and tubes with a sphere with tubes [H4] along two oppositely oriented tubes.

Disks without interior punctures are closed under boundary sewing operations. This closed structure is nothing but the operad of disks with strips, which was introduced and studied in [HKo1] and denoted as Υ . In [HKo1], Huang and the author introduced the notion of open-string vertex operator algebra and showed that open-string vertex operator algebras of central charge c are algebras over $\hat{\Upsilon}^c$, which is a $\mathbb{C}$ -extension of Υ .

The moduli space of spheres with tubes equipped with sewing operations has a structure of partial operad, called sphere partial operad and denoted as K . It is well understood by the works of Huang [H1][H2][H4]. Similar to the study of Υ , we can study $\mathfrak{S}$ in the framework of sphere partial operad K by embedding $\mathfrak{S}$ into K via a

doubling map δ [A][C1][HKo1]. The interior sewing operations on $\mathfrak{S}$ correspond to a double-sewing “action” of K on the image of δ (see (2.23)). As a consequence, the closed theory in an open-closed conformal field theory must contain both chiral part and anti-chiral part. Such closed theories were studied in [HKo2][K2][HKo3] in terms of the so-called (conformal) full field algebra and variants of it. More precisely, we showed in [K2] that conformal full field algebras over $V^L \otimes V^R$, where V^L and V^R are vertex operator algebras of central charge c^L and c^R respectively and satisfy certain finiteness and reductivity conditions, are algebras over the partial operad $\tilde{K}^{c^L} \otimes \overline{\tilde{K}^{c^R}}$, where $\tilde{K}^{c^L}$ and $\overline{\tilde{K}^{c^R}}$ are line bundles over the base space K [HKo1][K2].

We are interested in algebras over $\tilde{\mathfrak{S}}^c$, which is a $\mathbb{C}$ -extension of $\mathfrak{S}$. There is a natural action of $\tilde{K}^c \otimes \overline{\tilde{K}^c}$ on $\tilde{\mathfrak{S}}^c$ induced by interior sewing operations [HKo1]. Therefore, it is natural to expect that an algebra over $\tilde{\mathfrak{S}}^c$ with certain natural properties should contain a conformal full field algebra V_{cl} and an open-string vertex operator algebra V_{op} . Moreover, V_{cl} should “act” on V_{op} according to the action of $\tilde{K}^c \otimes \overline{\tilde{K}^c}$ on $\tilde{\mathfrak{S}}^c$. This requires certain compatibility between V_{cl} and V_{op} . One of the compatibility conditions is the so-called conformal invariant boundary condition. It roughly means that the two vertex operator algebras generated by the left and right Virasoro elements in V_{cl} must match in some way with that generated from the Virasoro element in V_{op} . In this case, we also call it a boundary condition preserving conformal symmetry. In this work, we only study those boundary conditions which preserve an enlarged symmetry given by a vertex operator algebra U (see [FS1][FS2] for general situations). Such structure is formalized by a notion called open-closed field algebra over U (see Definition 1.25). When U satisfies conditions in Theorem 0.1, we show that an open-closed field algebra over U canonically gives an algebra over $\tilde{\mathfrak{S}}^c$.

Open-closed field algebras over U are difficult to study for general U . The following Theorem proved by Huang in [H8] is important for us to simplify the situation.

Theorem 0.1. *Let V be a vertex operator algebra with central charge c satisfying the following conditions:*

1. *Every $\mathbb{C}$ -graded generalized V -module is a direct sum of $\mathbb{C}$ -graded irreducible V -modules,*
2. *There are only finitely many inequivalent $\mathbb{C}$ -graded irreducible V -modules,*
3. *Every $\mathbb{R}$ -graded irreducible V -module satisfies the C_1 -cofiniteness condition.*

Then the direct sum of all (in-equivalent) irreducible V -modules has a natural structure of intertwining operator algebra and the category of V -modules, denoted as $\mathcal{C}_V$, has a natural structure of vertex tensor category. In particular, $\mathcal{C}_V$ has natural structure of braided tensor category.

Assumption 0.2. In this work, we fix a vertex operator algebra V , which is assumed to satisfy the conditions in Theorem 0.1 without further announcement.

The notion of open-closed field algebra over V can be described by very few data and axioms in the framework of intertwining operator algebra (see Theorem 1.28 for a precise statement). Moreover, such algebra has a very simple categorical formulation. We will introduce a tensor-categorical notion called open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebra (see Definition 3.13) and show that the category of open-closed field algebras over V is isomorphic to the

category of open-closed $\mathcal{C}_V|\mathcal{C}_{V\otimes V}$ -algebras. The categorical formulation of open-closed field algebra over V makes it very easy to construct examples.

One of important developments on open-closed conformal field theories recently is a series of works [FFFS][FS3][FRS1]-[FRS4] [FjFRS] by Felder, Fröhlich, Fuchs, Runkel, Schweigert, Fjelstad, on rational open-closed conformal field theories using the theories of modular tensor category and 3-dimensional topological field theories. Assuming the existence of the structure of a modular tensor category on the category of modules for a vertex operator algebra and the existence of conformal blocks with monodromies compatible with the modular tensor category and all the necessary convergence properties, they constructed conformal blocks for open-closed conformal field theories of all genus and proved their factorization properties and invariance properties under the actions of mapping class groups. Our approach [HKo1]-[HKo3][K2][K3] is somewhat complement to theirs. We leave a detailed discussion of the relation between these two approaches to future publications.

The layout of this work is as follows. In Section 1, we review some old notions such as open-string vertex (operator) algebra, (conformal) full field algebra and variants of them. We also introduce the notions of open-closed field algebra and open-closed field algebra over U , and study their basic properties. In Section 2, we give an operadic formulation of open-closed field algebra over V . In particular, we show that an open-closed field algebra over V canonically gives an algebra over $\tilde{\mathfrak{S}}^c$. In Section 3, we give a categorical formulation of the notion of open-closed field algebra over V .

Convention of notations: $\mathbb{N}, \mathbb{Z}, \mathbb{Z}_+, \mathbb{R}, \mathbb{R}_+, \mathbb{C}$ denote the set of natural numbers, integers, positive integers, real numbers, positive real numbers, complex numbers, respectively. Let $\mathbb{H} = \{z \in \mathbb{C} | \text{Im} z > 0\}$, $\overline{\mathbb{H}} = \{z \in \mathbb{C} | \text{Im} z < 0\}$, $\tilde{\mathbb{H}} = \mathbb{H} \cup \mathbb{R}$, $\tilde{\overline{\mathbb{H}}} = \overline{\mathbb{H}} \cup \mathbb{R}$. We use $\hat{\mathbb{R}}, \hat{\mathbb{H}}, \hat{\tilde{\mathbb{H}}}$ to denote the one point compactification of $\mathbb{R}, \mathbb{H}, \tilde{\mathbb{H}}$ respectively. The following notations will also be used: $\forall n \in \mathbb{Z}_+$,

$$\Lambda^n := \{(r_1, \dots, r_n) \in \mathbb{R}^n | r_1 > \dots > r_n \geq 0\}, \quad (0.1)$$

$$M_{\mathbb{H}}^n := \{(z_1, \dots, z_n) \in \mathbb{H}^n | z_i \neq z_j, \text{ for } i, j = 1, \dots, n \text{ and } i \neq j\}, \quad (0.2)$$

$$M_{\mathbb{C}}^n := \{(z_1, \dots, z_n) \in \mathbb{C}^n | z_i \neq z_j, \text{ for } i, j = 1, \dots, n \text{ and } i \neq j\}. \quad (0.3)$$

The ground field is always assumed to be $\mathbb{C}$.

Acknowledgment This work grows from a chapter in author's thesis. I want to thank my advisor Yi-Zhi Huang for introducing this interesting field to me, for his many valuable suggestions for improvement and for his constant support. I also want to thank J. Lepowsky, J. Fuchs and C. Schweigert for many inspiring conversations, and I. Runkel for kindly sending me his thesis and one of his computer programs.

1 Open-closed field algebras

Let G be an abelian group. For any G -graded vector space $F = \bigoplus_{n \in G} F_{(n)}$ and any $n \in G$, we shall use P_n to denote the projection from F or $\overline{F} = \prod_{n \in G} F_{(n)}$ to $F_{(n)}$. We give F and its graded dual $F' = \bigoplus_{n \in G} F_{(n)}^*$ the topology induced from the pairing between F and F' . We also give $\text{Hom}(F, \overline{F})$ the topology induced from the linear functionals on $\text{Hom}(F, \overline{F})$ given by $f \mapsto \langle v', f(v) \rangle$ for $f \in \text{Hom}(F, \overline{F})$, $v \in F$ and $v' \in F'$. For any vector space F and any set S , $F^{\otimes 0} = \mathbb{C}$ and S^0 is an one-point set $\{*\}$.

1.1 Boundary field algebras and open-string vertex algebras

Definition 1.1. A *boundary field algebra* is a $\mathbb{R}$ -graded vector space V_{op} with grading operator being $\mathbf{d}_{op}$, together with a correlation-function map for each $n \in \mathbb{N}$:

$$\begin{aligned} m_{op}^{(n)} : V_{op}^{\otimes n} \times \Lambda^n &\rightarrow \overline{V}_{op} \\ (u_1 \otimes \cdots \otimes u_n, (r_1, \dots, r_n)) &\mapsto m_{op}^{(n)}(u_1, \dots, u_n; r_1, \dots, r_n) \end{aligned}$$

and an operator $D_{op} \in \text{End } V_{op}$, satisfying the following axioms:

1. For each $n \in \mathbb{Z}_+$, $m_{op}^{(n)}(u_1, \dots, u_n; r_1, \dots, r_n)$ is linear in $u_1, \dots, u_n$ and smooth in $r_1, \dots, r_n$.
2. $\forall u \in V_{op}$, $m_{op}^{(1)}(u; 0) = u$ and $\mathbf{1}_{op} := m_{op}^{(0)}(1) \in (V_{op})_{(0)}$.
3. *Convergence property:* For $n \in \mathbb{Z}_+$, $k \in \mathbb{N}$ and $i = 1, \dots, n$, $u_1, \dots, u_n, u_1^{(i)}, \dots, u_k^{(i)} \in V_{op}$, the following series¹

$$\sum_{s \in \mathbb{R}} m_{op}^{(n)}(u_1, \dots, u_{i-1}, P_s m_{op}^k(u_1^{(1)}, \dots, u_k^{(i)}, r_1^{(1)}, \dots, r_k^{(i)}), u_{i+1}, \dots, u_n; r_1, \dots, r_n) \quad (1.1)$$

converges absolutely, whenever $r_1^{(i)} < |r_i - r_j|$ for all $j \neq i$, to

$$\begin{aligned} m_{op}^{(n+k-1)}(u_1, \dots, u_{i-1}, u_1^{(i)}, \dots, u_k^{(i)}, u_{i+1}, \dots, u_n; \\ r_1, \dots, r_{i-1}, r_i + r_1^{(i)}, \dots, r_i + r_k^{(i)}, r_{i+1}, \dots, r_n). \end{aligned} \quad (1.2)$$

4. *$\mathbf{d}_{op}$ -bracket property:*

$$e^{\mathbf{ad}_{op}} m_{op}^{(n)}(u_1, \dots, u_n; r_1, \dots, r_n) = m_{op}^{(n)}(e^{\mathbf{ad}_{op}} u_1, \dots, e^{\mathbf{ad}_{op}} u_n; e^a r_1, \dots, e^a r_n). \quad (1.3)$$

for $n \in \mathbb{Z}_+$, $r_1 > \cdots > r_n \geq 0$, $r \in \mathbb{R}$ and $u_1, \dots, u_n \in V_{op}$.

5. *D_{op} -property:* For $u_1, \dots, u_n \in V_{op}$, $r_1 > \cdots > r_n \geq 0$ and $r_n + a \geq 0$,

$$e^{aD_{op}} m_{op}^{(n)}(u_1, \dots, u_n; r_1, \dots, r_n) = m_{op}^{(n)}(u_1, \dots, u_n, r_1 + a, \dots, r_n + a). \quad (1.4)$$

We denote such a boundary field algebra as $(V_{op}, m_{op}, \mathbf{d}_{op}, D_{op})$. *Homomorphisms*, *isomorphisms* and *subalgebras* of boundary field algebra are defined in the obvious way.

Let the map $Y_{op} : V_{op}^{\otimes(2)} \times \mathbb{R}_+ \rightarrow \overline{V}_{op}$ be defined by

$$Y_{op} : (u \otimes v, r) \mapsto Y_{op}(u, r)v = m_2(u, v; r, 0). \quad (1.5)$$

Then by the convergence property, we have

$$Y_{op}(\mathbf{1}_{op}, r) = \text{id}_F, \quad (1.6)$$

$$\lim_{r \rightarrow 0} Y_{op}(u, r)\mathbf{1}_{op} = u, \quad (1.7)$$

¹That the number of nonvanishing terms in the sum is countable is automatically assumed.

for $u \in V_{op}$. (1.7) implies that the map $u \mapsto Y_{op}(u, r)$ is one-to-one. By (1.3), we have

$$e^{aD_{op}} Y_{op}(u, r) e^{-aD_{op}} = Y_{op}(e^{aD_{op}} u, ar) \quad (1.8)$$

for $u \in V_{op}, r \in \mathbb{R}_+$ and $a \in \mathbb{R}$. By (1.4), we also have

$$e^{aD_{op}} Y_{op}(u, r) e^{-aD_{op}} = Y_{op}(u, r + a) \quad (1.9)$$

for $u \in V_{op}, r > 0, r + a > 0$. Moreover, using (1.7) and (1.4), we obtain the identity:

$$Y_{op}(v, r) \mathbf{1}_{op} = m_{op}^{(1)}(v; r) = e^{rD_{op}} v. \quad (1.10)$$

If we assume some analytic properties on $m_{op}^{(n)}$, it is possible to use Y_{op} to generate all $m_{op}^{(n)}$. This motivate us to introduce the notion of open-string vertex algebra in [HKo1]. The definition given here is a refinement of that in [HKo1].

Definition 1.2. An *open-string vertex algebra* is an $\mathbb{R}$ -graded vector space $V_{op} = \bigoplus_{n \in \mathbb{R}} (V_{op})_{(n)}$ (graded by *weights*) equipped with a *vertex map*:

$$\begin{aligned} Y_{op} : (V_{op} \otimes V_{op}) \times \mathbb{R}_+ &\rightarrow \overline{V}_{op} \\ (u \otimes v, r) &\mapsto Y_{op}(u, r)v, \end{aligned} \quad (1.11)$$

a *vacuum* $\mathbf{1}_{op} \in V_{op}$ and an operator $D_{op} \in \text{End } V_{op}$ of weight 1, satisfying the following conditions:

1. *Vertex map weight property*: For $s_1, s_2 \in \mathbb{R}$, there exists a finite subset $S(s_1, s_2) \subset \mathbb{R}$ such that the image of $(\bigoplus_{s \in s_1 + \mathbb{Z}} (V_{op})_{(s)}) \otimes (\bigoplus_{s \in s_2 + \mathbb{Z}} (V_{op})_{(s)})$ under the vertex map Y_{op} is in $\prod_{s \in S(s_1, s_2) + \mathbb{Z}} (V_{op})_{(s)}$.
2. *Vacuum properties*:
 - (a) *identity property*: For any $r \in \mathbb{R}_+$, $Y_{op}(\mathbf{1}_{op}, r) = \text{id}_{V_{op}}$,
 - (b) *creation property*: $\forall u \in V_{op}, \lim_{r \rightarrow 0} Y_{op}(u, r) \mathbf{1}_{op} = u$.
3. *Convergence properties*:

For $n \in \mathbb{N}$, $u_1, \dots, u_n, v \in V_{op}$ and $v' \in V'_{op}$, the series

$$\begin{aligned} &\langle v', Y_{op}(u_1, r_1) \dots Y_{op}(u_n, r_n) v \rangle \\ &= \sum_{m_1, \dots, m_{n-1}} \langle v', Y_{op}(u_1, r_1) P_{m_1} Y_{op}(u_2, r_2) \dots P_{m_{n-1}} Y_{op}(u_n, r_n) v \rangle \end{aligned}$$

converges absolutely when $r_1 > \dots > r_n > 0$ and is a restriction to the domain $\{r_1 > \dots > r_n > 0\}$ of an (possibly multivalued) analytic function in $(\mathbb{C}^\times)^n$ with only possible singularities at $r_i = r_j$ for $1 \leq i, j \leq n$ and $i \neq j$.

- (a) For $u_1, u_2, v \in V_{op}, v' \in V'_{op}$, the series

$$\langle v', Y_{op}(Y_{op}(u_1, r_0) u_2, r_2) v \rangle = \sum_m \langle v', Y_{op}(P_m Y_{op}(u_1, r_0) u_2, r_2) v \rangle$$

converges absolutely when $r_2 > r_0 > 0$.

4. *Associativity*: For $u_1, u_2, v \in V_{op}$ and $v' \in V'_{op}$,

$$\langle v', Y_{op}(u_1, r_1)Y_{op}(u_2, r_2)v \rangle = \langle v', Y_{op}(Y_{op}(u_1, r_1 - r_2)u_2, r_2)v \rangle$$

for $r_1, r_2 \in \mathbb{R}$ satisfying $r_1 > r_2 > r_1 - r_2 > 0$.

5. *$\mathbf{d}_{op}$ -bracket property*: Let $\mathbf{d}_{op}$ be the grading operator on V_{op} . For $u \in V_{op}$ and $r \in \mathbb{R}_+$,

$$[\mathbf{d}_{op}, Y_{op}(u, r)] = Y_{op}(\mathbf{d}_{op}u, r) + r \frac{d}{dr} Y_{op}(u, r). \quad (1.12)$$

6. *D_{op} -derivative property*: We still use D_{op} to denote the natural extension of D_{op} to $\text{Hom}(\overline{V}_{op}, \overline{V}_{op})$. For $u \in V_{op}$, $Y_{op}(u, r)$ as a map from $\mathbb{R}_+$ to $\text{Hom}(V_{op}, \overline{V}_{op})$ is differentiable and

$$\frac{d}{dr} Y_{op}(u, r) = [D_{op}, Y_{op}(u, r)] = Y_{op}(D_{op}u, r). \quad (1.13)$$

Homomorphisms, isomorphisms, subalgebras of open-string vertex algebras are defined in the obvious way.

We denote such algebra by $(V_{op}, Y_{op}, \mathbf{1}_{op}, \mathbf{d}_{op}, D_{op})$ or simply V_{op} . For $u \in V_{op}$, it was shown in [HKo1] that there is a formal vertex operator

$$Y_{op}^f(u, x) = \sum_{n \in \mathbb{R}} u_n x^{-n-1}, \quad (1.14)$$

where $u_n \in \text{End } V_{op}$, so that

$$Y_{op}^f(u, x)|_{x=r} = Y_{op}(u, r). \quad (1.15)$$

We can also replace x by complex variable z if we choose a branch cut. For any $z \in \mathbb{C}^\times$ and $n \in \mathbb{R}$, we define

$$\log z := \log |z| + \arg z, 0 \leq \arg z < 2\pi. \quad (1.16)$$

But for power functions, we distinguish two types of complex variables, z (or $z_1, z_2, \dots$) and $\bar{z}_1, \zeta$ (or $\bar{z}_1, \zeta_1, \bar{z}_2, \zeta_2, \dots$). We define

$$z^n := e^{n \log z}, \quad \bar{z}^n := e^{n \overline{\log z}}, \quad \zeta^n := e^{n \overline{\log \zeta}}. \quad (1.17)$$

Proposition 1.3. *An open-string vertex algebra canonically gives a boundary field algebra.*

Proof. The proof is standard. We omit it here. ■

Definition 1.4. An *open-string vertex operator algebra* is an open-string vertex algebra, together with a conformal element ω_{op} , satisfying the following conditions:

7. *grading-restriction conditions*: For all $n \in \mathbb{R}$, $\dim(V_{op})_{(n)} < \infty$ and $(V_{op})_{(n)} = 0$ when n is sufficiently negative.

8. *Virasoro relations*: The vertex operator associated to ω_{op} has the following expansion:

$$Y_{op}(\omega_{op}, r) = \sum_{n \in \mathbb{Z}} L(n) r^{-n-2}$$

where $L(n)$ satisfying the following condition: $\forall m, n \in \mathbb{Z}$,

$$[L(m), L(n)] = (m - n)L(m + n) - \frac{c}{12}(m^3 - m)\delta_{m+n,0},$$

for some $c \in \mathbb{C}$.

9. *commutator formula for Virasoro operators and formal vertex operators (or component operators)*: For $v \in V_{op}$, $Y_{op}^f(\omega_{op}, x)v$ involves only finitely many negative powers of x and

$$[Y_{op}^f(\omega_{op}, x_1), Y_{op}^f(v, x_2)] = \text{Res}_{x_0} x_2^{-1} \delta\left(\frac{x_1 - x_0}{x_2}\right) Y_{op}^f(Y_{op}^f(\omega_{op}, x_0)v, x_2).$$

10. *$L(0)$ -grading property and $L(-1)$ -derivative property*: $L(0) = \mathbf{d}_{op}$, $L(-1) = D_{op}$.

We shall denote the open-string vertex operator algebra defined above by $(V_{op}, Y_{op}, \mathbf{1}_{op}, \omega_{op})$ or simply V_{op} . The complex number c in the definition is called *central charge*.

The *meromorphic center* of V_{op} is defined by

$$C_0(V_{op}) = \left\{ u \in \oplus_{n \in \mathbb{Z}} (V_{op})_{(n)} \mid Y_{op}^f(u, x) \in (\text{End } V_{op})[[x, x^{-1}]], \right. \\ \left. Y_{op}^f(v, x)u = e^{xD} Y_{op}^f(u, -x)v, \forall v \in V_{op} \right\}.$$

It was shown in [HKo1] that the meromorphic center of a grading-restricted open-string vertex (operator) algebra is a grading-restricted vertex (operator) algebra.

Let U be a grading-restricted vertex (operator) algebra. If there is a monomorphism $\iota_{op} : U \hookrightarrow C_0(V_{op})$ of grading-restricted vertex (operator) algebra, we call V_{op} an open-string vertex (operator) algebra over U , and denote it by $(V_{op}, Y_{op}, \iota_{op})$ or simply by V_{op} . In this case, the formal vertex operator Y_{op}^f is an intertwining operator of type $\binom{V_{op}}{V_{op} V_{op}}$, where V_{op} is a U -module.

Remark 1.5. Open-string vertex operator algebra can be viewed as a noncommutative generalization of vertex operator algebra. Other noncommutative generalizations of vertex (operator) algebra were also studied in the literature [FR][B][EK][BKa][L1][L2][L3].

1.2 Full field algebras

Definition 1.6. A $\mathbb{R} \times \mathbb{R}$ -graded *full field algebra* is an $\mathbb{R} \times \mathbb{R}$ -graded vector space $V_{cl} = \oplus_{m,n \in \mathbb{R}} (V_{cl})_{(m,n)}$ (graded by *left weight* wt^L and *right weight* wt^R with left and right grading operators $\mathbf{d}^L$ and $\mathbf{d}^R$), equipped with *correlation-function maps*

$$m_{cl}^{(n)} : V_{cl}^{\otimes n} \times M_{\mathbb{C}}^n \rightarrow \overline{V}_{cl} \\ (u_1 \otimes \cdots \otimes u_n, (z_1, \dots, z_n)) \mapsto m_{cl}^{(n)}(u_1, \dots, u_n; z_1, \bar{z}_1, \dots, z_n, \bar{z}_n),$$

for each $n \in \mathbb{N}$, and operators D^L and D^R of weights $(1, 0)$ and $(0, 1)$ respectively, satisfying the following axioms:

1. *Single-valuedness property:* $e^{2\pi i(\mathbf{d}^L - \mathbf{d}^R)} = \text{id}_{V_{cl}}$.
2. For $n \in \mathbb{Z}_+$, $m_{cl}^{(n)}(u_1, \dots, u_n; z_1, \bar{z}_1, \dots, z_n, \bar{z}_n)$ is linear in $u_1, \dots, u_n$ and smooth in the real and imaginary parts of $z_1, \dots, z_n$.
3. *Identity properties:* $\forall u \in V_{cl}$, $m_1(u; 0, 0) = u$ and $\mathbf{1}_{cl} := m_{cl}^{(0)}(1) \in (V_{cl})_{(0,0)}$.
4. *Convergence property:* For $k \in \mathbb{Z}_+$ and $u_1, \dots, u_n, u_1^{(i)}, \dots, u_k^{(i)} \in V_{cl}$ and $i = 1, \dots, n$, the series

$$\sum_{p,q \in \mathbb{R}} m_{cl}^{(n)}(u_1, \dots, u_{i-1}, P_{(p,q)} m_{cl}^{(k)}(v_1, \dots, v_k; z_1^{(i)}, \bar{z}_1^{(i)}, \dots, z_k^{(i)}, \bar{z}_k^{(i)}), \\ u_{i+1}, \dots, u_n; z_1, \bar{z}_1, \dots, z_n, \bar{z}_n) \quad (1.18)$$

converges absolutely to

$$m_{cl}^{(n+k-1)}(u_1, \dots, u_{i-1}, v_1, \dots, v_k, u_{i+1}, \dots, u_n; z_1, \bar{z}_1, \dots, z_{i-1}, \bar{z}_{i-1}, \\ z_i + z_1^{(i)}, \bar{z}_i + \bar{z}_1^{(i)} \dots, z_i + z_k^{(i)}, \bar{z}_i + \bar{z}_k^{(i)}, z_{i+1}, \bar{z}_{i+1}, \dots, z_n, \bar{z}_n) \quad (1.19)$$

whenever $|z_p^{(i)}| < |z_i - z_j|$ for all $j = 1, \dots, n$, $i \neq j$ and for $p = 1, \dots, k$.

5. *Permutation property:* For $n \in \mathbb{Z}_+$ and $\sigma \in S_n$, we have

$$m_{cl}^{(n)}(u_1, \dots, u_n; z_1, \bar{z}_1, \dots, z_n, \bar{z}_n) \\ = m_{cl}^{(n)}(u_{\sigma(1)}, \dots, u_{\sigma(n)}; z_{\sigma(1)}, \bar{z}_{\sigma(1)}, \dots, z_{\sigma(n)}, \bar{z}_{\sigma(n)}) \quad (1.20)$$

for $u_1, \dots, u_n \in V_{cl}$ and $(z_1, \dots, z_n) \in M_{\mathbb{C}}^n$.

6. $\mathbf{d}^L$ and $\mathbf{d}^R$ property: For $u_1, \dots, u_n \in V_{cl}$ and $a \in \mathbb{C}$,

$$e^{a\mathbf{d}^L} e^{\bar{a}\mathbf{d}^R} m_{cl}^{(n)}(u_1, \dots, u_n; z_1, \bar{z}_1, \dots, z_n, \bar{z}_n) \\ = m_{cl}^{(n)}(e^{a\mathbf{d}^L} e^{\bar{a}\mathbf{d}^R} u_1, \dots, e^{a\mathbf{d}^L} e^{\bar{a}\mathbf{d}^R} u_n, e^a z_1, e^{\bar{a}} \bar{z}_1, \dots, e^a z_n, e^{\bar{a}} \bar{z}_n). \quad (1.21)$$

7. D^L and D^R property: $[D^L, D^R] = 0$ and for $u_1, \dots, u_n \in V_{cl}$ and $a \in \mathbb{C}$,

$$e^{aD^L} e^{\bar{a}D^R} m_{cl}^{(n)}(u_1, \dots, u_n; z_1, \bar{z}_1, \dots, z_n, \bar{z}_n) \\ = m_{cl}^{(n)}(u_1, \dots, u_n, z_1 + a, \bar{z}_1 + \bar{a}, \dots, z_n + a, \bar{z}_n + \bar{a}). \quad (1.22)$$

We denote the $\mathbb{R} \times \mathbb{R}$ -graded full field algebra defined above by

$$(V_{cl}, m_{cl}, \mathbf{d}^L, \mathbf{d}^R, D^L, D^R)$$

or simply by V_{cl} .

Subalgebra, homomorphism, monomorphism, epimorphism and isomorphism of full field algebra can be naturally defined.

Let $\mathbb{Y} : V_{cl}^{\otimes 2} \times \mathbb{C}^\times \rightarrow \overline{V}_{cl}$ be so that

$$\mathbb{Y} : (u \otimes v, z) \mapsto \mathbb{Y}(u; z, \bar{z})v := m_2(u \otimes v; z, \bar{z}, 0, 0). \quad (1.23)$$

Then by the convergence property, it is easy to see that

$$\mathbb{Y}(\mathbf{1}_{cl}; z, \bar{z}) = \text{id}_{V_{cl}}, \quad (1.24)$$

$$\lim_{z \rightarrow 0} \mathbb{Y}(u; z, \bar{z}) \mathbf{1}_{cl} = u, \quad \forall u \in V_{cl}. \quad (1.25)$$

Moreover, it is also not hard to show the following two properties of $\mathbb{Y}$:

1. The $\mathbf{d}^L$ - and $\mathbf{d}^R$ -bracket properties:

$$[\mathbf{d}^L, \mathbb{Y}(u; z, \bar{z})] = z \frac{\partial}{\partial z} \mathbb{Y}(u; z, \bar{z}) + \mathbb{Y}(\mathbf{d}^L u; z, \bar{z}), \quad (1.26)$$

$$[\mathbf{d}^R, \mathbb{Y}(u; z, \bar{z})] = \bar{z} \frac{\partial}{\partial \bar{z}} \mathbb{Y}(u; z, \bar{z}) + \mathbb{Y}(\mathbf{d}^R u; z, \bar{z}). \quad (1.27)$$

2. The D^L - and D^R -derivative property:

$$[D^L, \mathbb{Y}(u; z, \bar{z})] = \mathbb{Y}(D^L u; z, \bar{z}) = \frac{\partial}{\partial z} \mathbb{Y}(u; z, \bar{z}), \quad (1.28)$$

$$[D^R, \mathbb{Y}(u; z, \bar{z})] = \mathbb{Y}(D^R u; z, \bar{z}) = \frac{\partial}{\partial \bar{z}} \mathbb{Y}(u; z, \bar{z}). \quad (1.29)$$

It was shown in [HKo2] that we have the following expansion:

$$\mathbb{Y}(u; z, \bar{z}) = \sum_{r, s \in \mathbb{R}} \mathbb{Y}_{l,r}(u) e^{(-l-1) \log z} e^{(-r-1) \overline{\log z}} \quad (1.30)$$

where $\mathbb{Y}_{l,r}(u) \in \text{End } F$ with $\text{wt}^L \mathbb{Y}_{l,r}(u) = \text{wt}^L u - l - 1$ and $\text{wt}^R \mathbb{Y}_{l,r}(u) = \text{wt}^R u - r - 1$. Moreover, the expansion above is unique. Let x and $\bar{x}$ be independent and commuting formal variables. We define the *formal full vertex operator* $\mathbb{Y}_f$ associated to $u \in V_{cl}$ by

$$\mathbb{Y}_f(u; x, \bar{x}) = \sum_{l, r \in \mathbb{R}} \mathbb{Y}_{l,r}(u) x^{-l-1} \bar{x}^{-r-1}. \quad (1.31)$$

For nonzero complex numbers z and ζ , we can substitute $e^{r \log z}$ and $e^{s \overline{\log \zeta}}$ for x^r and $\bar{x}^s$ respectively in $\mathbb{Y}_f(u; x, \bar{x})$ to obtain an operator

$$\mathbb{Y}_{\text{an}}(u; z, \zeta) : V_{cl} \times (\mathbb{C}^\times)^2 \rightarrow \overline{V}_{cl}$$

called *analytic full vertex operator*.

Definition 1.7. An $\mathbb{R} \times \mathbb{R}$ -graded full field algebra $(V_{cl}, m_{cl}, \mathbf{d}^L, \mathbf{d}^R, D^L, D^R)$ is called *grading-restricted* if it satisfies the following grading-restriction conditions:

1. There exists $M \in \mathbb{R}$ such that $(V_{cl})_{(m,n)} = 0$ if $n < M$ or $m < M$.
2. $\dim(V_{cl})_{(m,n)} < \infty$ for $m, n \in \mathbb{R}$.

We say that V_{cl} is *lower-truncated* if V_{cl} satisfies the first grading restriction condition.

In this case, for $u \in V_{cl}$ and $k \in \mathbb{R}$, we have

$$\sum_{l+r=k} \mathbb{Y}_{l,r}(u) \in \text{End } V_{cl}$$

with total weight $\text{wt } u - k - 2$. We denote $\sum_{l+r=k} \mathbb{Y}_{l,r}(u)$ by $\mathbb{Y}_{k-1}(u)$. Then we have the expansion

$$\mathbb{Y}_f(u; x, x) = \sum_{k \in \mathbb{R}} \mathbb{Y}_k(u) x^{-k-1}, \quad (1.32)$$

where $\text{wt } \mathbb{Y}_k(u) = \text{wt } u - k - 1$. For given $u, v \in v_{cl}$, we have $\mathbb{Y}_k(u)w = 0$ for sufficiently large k .

Let $(V^L, Y^L, \mathbf{1}^L, \omega^L)$ and $(V^R, Y^R, \mathbf{1}^R, \omega^R)$ be vertex operator algebras. It was pointed out in [HKo2] that $V^L \otimes V^R$ has a natural structure of full field algebra. Let ι_{cl} be an injective homomorphism from the full field algebra $V^L \otimes V^R$ to V_{cl} . Then we have $\mathbf{1}_{cl} = \iota_{cl}(\mathbf{1}^L \otimes \mathbf{1}^R)$, $\mathbf{d}^L \circ \iota_{cl} = \iota_{cl} \circ (L^L(0) \otimes I_{V^R})$, $\mathbf{d}^R \circ \iota_{cl} = \iota_{cl} \circ (I_{V^L} \otimes L^R(0))$, $D^L \circ \iota_{cl} = \iota_{cl} \circ (L^L(-1) \otimes I_{V^R})$ and $D^R \circ \iota_{cl} = \iota_{cl} \circ (I_{V^L} \otimes L^R(-1))$. Moreover, V_{cl} has a *left conformal element* $\iota_{cl}(\omega^L \otimes \mathbf{1}^R)$ and an *right conformal element* $\iota_{cl}(\mathbf{1}^L \otimes \omega^R)$. We have the following operators on V_{cl} :

$$\begin{aligned} L^L(0) &= \text{Res}_x \text{Res}_{\bar{x}} \bar{x}^{-1} \mathbb{Y}_f(\iota_{cl}(\omega^L \otimes \mathbf{1}^R); x, \bar{x}), \\ L^R(0) &= \text{Res}_x \text{Res}_{\bar{x}} x^{-1} \mathbb{Y}_f(\iota_{cl}(\mathbf{1}^L \otimes \omega^R); x, \bar{x}), \\ L^L(-1) &= \text{Res}_x \text{Res}_{\bar{x}} \bar{x} \mathbb{Y}_f(\iota_{cl}(\omega^L \otimes \mathbf{1}^R); x, \bar{x}), \\ L^R(-1) &= \text{Res}_x \text{Res}_{\bar{x}} x \mathbb{Y}_f(\iota_{cl}(\mathbf{1}^L \otimes \omega^R); x, \bar{x}). \end{aligned}$$

Since these operators are operators on V_{cl} , it should be easy to distinguish them from those operators with the same notation but acting on V^L or V^R .

Definition 1.8. Let $(V^L, Y^L, \mathbf{1}^L, \omega^L)$ and $(V^R, Y^R, \mathbf{1}^R, \omega^R)$ be vertex operator algebras. A *full field algebra over $V^L \otimes V^R$* is a grading-restricted $\mathbb{R} \times \mathbb{R}$ -graded full field algebra $(V_{cl}, m_{cl}, \mathbf{d}^L, \mathbf{d}^R, D^L, D^R)$ equipped with an injective homomorphism ι_{cl} from the full field algebra $V^L \otimes V^R$ to V_{cl} such that $\mathbf{d}^L = L^L(0)$, $\mathbf{d}^R = L^R(0)$, $D^L = L^L(-1)$ and $D^R = L^R(-1)$. A full field algebra over $V^L \otimes V^R$ equipped with left and right conformal elements $\iota_{cl}(\omega^L \otimes \mathbf{1}^R)$ and $\iota_{cl}(\mathbf{1}^L \otimes \omega^R)$ is called *conformal full field algebra over $V^L \otimes V^R$* .

We shall denote the (conformal) full field algebra over $V^L \otimes V^R$ defined above by $(V_{cl}, m_{cl}, \iota_{cl})$ or simply by V_{cl} . From now on, we will not distinguish V^L with $V^L \otimes \mathbf{1}^R$ and $\iota_{cl}(V^L \otimes \mathbf{1}^R)$. Similarly for V^R with $\mathbf{1}^L \otimes V^R$ and $\iota_{cl}(\mathbf{1}^L \otimes V^R)$. For $u^L \in V^L$, $\mathbb{Y}_{an}(u^L; z, \zeta)$ is independent of ζ . So we simply denote it as $\mathbb{Y}_{an}(u^L, z)$. Similarly, we denote $\mathbb{Y}_{an}(u^R; z, \zeta)$ as $\mathbb{Y}_{an}(u^R, \zeta)$ for $u^R \in V^R$.

1.3 Open-closed field algebras

Definition 1.9. An open-closed field algebra consists of a $\mathbb{R} \times \mathbb{R}$ -graded full field algebra

$$(V_{cl}, m_{cl}, \mathbf{d}_{cl}^L, \mathbf{d}_{cl}^R, D_{cl}^L, D_{cl}^R)$$

and a $\mathbb{R}$ -graded vector space V_{op} with the grading operator $\mathbf{d}_{op}$ and an additional operator $D_{op} \in \text{End } V_{op}$, together with a map for each pair of $n, l \in \mathbb{N}$:

$$\begin{aligned} m_{cl-op}^{(l;n)} : V_{cl}^{\otimes l} \otimes V_{op}^{\otimes n} \times M_{\mathbb{H}}^l \times \Lambda^n &\rightarrow \overline{V}_{op} \\ (u_1 \otimes \cdots \otimes u_l \otimes v_1 \otimes \cdots \otimes v_n, (z_1, \dots, z_l; r_1, \dots, r_n)) &\mapsto \\ m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n), \end{aligned}$$

satisfying the following axioms:

1. $m_{cl-op}^{(l;n)}(u_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n)$ is linear in $u_1, \dots, v_n$ and smooth in $r_1, \dots, r_n, z_1, \dots, z_l$.
2. *Identity properties:* $m_{cl-op}^{(0;1)}(v; 0) = v, \forall v \in V_{op}$ and $\mathbf{1}_{op} := m_{cl-op}^{(0;0)}(1) \in (V_{op})_{(0)}$.
3. *Convergence properties:*

- (a) For $u_1, \dots, u_l, \tilde{u}_1, \dots, \tilde{u}_k \in V_{cl}$, $v_1, \dots, v_n, \tilde{v}_1, \dots, \tilde{v}_m \in V_{op}$ and $i = 1, \dots, n$, the following series

$$\sum_{n_1 \in \mathbb{R}} m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_{i-1}, P_{n_1} m_{cl-op}^{(k;m)}(\tilde{u}_1, \dots, \tilde{u}_k; \tilde{v}_1, \dots, \tilde{v}_m; z_1^{(i)}, \overline{z_1^{(i)}}, \dots, z_k^{(i)}, \overline{z_k^{(i)}}; r_1^{(i)}, \dots, r_m^{(i)}); v_{i+1}, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \quad (1.33)$$

converges absolutely when $|z_s - r_i|, |r_t - r_i| > |z_p^{(i)}|, r_q^{(i)} \geq 0$ for all $s = 1, \dots, l$, $t = 1, \dots, n$, $t \neq i$, $p = 1, \dots, k$ and $q = 1, \dots, m$ to

$$m_{cl-op}^{(l+k;n+m-1)}(u_1, \dots, u_l, \tilde{u}_1, \dots, \tilde{u}_k; v_1, \dots, v_{i-1}, \tilde{v}_1, \dots, \tilde{v}_m, v_{i+1}, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l, z_1^{(i)}, \overline{z_1^{(i)}}, \dots, z_k^{(i)}, \overline{z_k^{(i)}}; r_1, \dots, r_{i-1}, r_i + r_1^{(i)}, \dots, r_i + r_m^{(i)}, r_{i+1}, \dots, r_n). \quad (1.34)$$

- (b) For $u_1, \dots, u_l, \tilde{u}_1, \dots, \tilde{u}_k \in V_{cl}$, $v_1, \dots, v_n \in V_{op}$ and $i = 1, \dots, n$, the following series

$$\sum_{n_1, n_2 \in \mathbb{R}} m_{cl-op}^{(l;n)}(u_1, \dots, u_{i-1}, P_{(n_1, n_2)} m_{cl}^{(k)}(\tilde{u}_1, \dots, \tilde{u}_k; z_1^{(i)}, \overline{z_1^{(i)}}, \dots, z_k^{(i)}, \overline{z_k^{(i)}}), u_{i+1}, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \quad (1.35)$$

converges absolutely, when $|z_s - z_i|, |r_t - z_i| > |z_p^{(i)}|$ for all $s = 1, \dots, l$, $s \neq i$, $t = 1, \dots, n$, $p = 1, \dots, k$, to

$$m_{cl-op}^{(l+k-1;n)}(u_1, \dots, u_{i-1}, \tilde{u}_1, \dots, \tilde{u}_k, u_{i+1}, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_{i-1}, \bar{z}_{i-1}, z_i + z_1^{(i)}, \overline{z_i + z_1^{(i)}}, \dots, z_i + z_k^{(i)}, \overline{z_i + z_k^{(i)}}, z_{i+1}, \bar{z}_{i+1}, \dots, z_l, \bar{z}_l; r_1, \dots, r_n). \quad (1.36)$$

4. *Permutation axiom*: For $u_1, \dots, u_l \in V_{cl}$, $v_1, \dots, v_n \in V_{op}$ and $\sigma \in S_l$,

$$\begin{aligned} & m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(u_{\sigma(1)}, \dots, u_{\sigma(l)}; v_1, \dots, v_n; z_{\sigma(1)}, \bar{z}_{\sigma(1)}, \dots, z_{\sigma(l)}, \bar{z}_{\sigma(l)}; r_1, \dots, r_n). \end{aligned} \quad (1.37)$$

5. $\mathbf{d}_{op}-$, $\mathbf{d}_{cl}^L-$ and $\mathbf{d}_{cl}^R$ -property: For $u_1, \dots, u_l \in V_{cl}$, $v_1, \dots, v_n \in V_{op}$ and $a \in \mathbb{R}$,

$$\begin{aligned} & e^{a\mathbf{d}_{op}} m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(e^{a(\mathbf{d}_{cl}^L + \mathbf{d}_{cl}^R)} u_1, \dots, e^{a(\mathbf{d}_{cl}^L + \mathbf{d}_{cl}^R)} u_l; e^{a\mathbf{d}_{op}} v_1, \dots, e^{a\mathbf{d}_{op}} v_n; e^a z_1, e^a \bar{z}_1, \dots, e^a z_l, e^a \bar{z}_l; e^a r_1, \dots, e^a r_n). \end{aligned} \quad (1.38)$$

6. D_{op} -property: For $u_1, \dots, u_l \in V_{cl}$, $v_1, \dots, v_n \in V_{op}$ and $r_n + a \geq 0$,

$$\begin{aligned} & e^{aD_{op}} m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1 + a, \bar{z}_1 + a, \dots, z_l + a, \bar{z}_l + a; r_1 + a, \dots, r_n + a). \end{aligned} \quad (1.39)$$

We denote such algebra by $(V_{cl}, V_{op}, m_{cl-op})$ for simplicity. *Homomorphisms, isomorphisms, subalgebras* of open-closed field algebras are defined in the obvious way.

Remark 1.10. From above definition, it is clear that an open-closed field algebra automatically includes a boundary field algebra as a substructure.

We discuss a few results which follow immediately from the definition. By the identity properties and (1.39), we have, for $a \geq 0$,

$$m_{cl-op}^{(0;1)}(v; a) = e^{aD_{op}} m_{cl-op}^{(0;1)}(v; 0) = e^{aD_{op}} v. \quad (1.40)$$

By (1.40) and the convergence property, for $i = 1, \dots, n$ and $a \in \mathbb{R}$, we obtain

$$\begin{aligned} & m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_{i-1}, e^{aD_{op}} v_i, v_{i+1}, \dots, v_n; \\ & \quad z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_{i-1}, m_{cl-op}^{(0;1)}(v_i; a), \\ & \quad v_{i+1}, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l \\ & \quad r_1, \dots, r_{i-1}, r_i + a, r_{i+1}, \dots, r_n) \end{aligned} \quad (1.41)$$

when $|r_j - r_i|, |z_k - r_i| > |a|$ for $j \neq i$ and $k = 1, \dots, l$. By (1.22) and the convergence property, we also have, for $j = 1, \dots, l$ and $b \in \mathbb{C}$,

$$\begin{aligned} & m_{cl-op}^{(l;n)}(u_1, \dots, u_{j-1}, e^{bD_{cl}^L + \bar{b}D_{cl}^R} u_j, u_{j+1}, \dots, u_l; v_1, \dots, v_n; \\ & \quad z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(u_1, \dots, u_{j-1}, m_{cl}^{(1)}(u_j; b, \bar{b}), u_{j+1}, \dots, u_l; \\ & \quad v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \\ &= m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; z_1, \bar{z}_1, \dots, z_{j-1}, \bar{z}_{j-1}, \\ & \quad z_j + b, \bar{z}_j + \bar{b}, z_{j+1}, \bar{z}_{j+1}, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) \end{aligned} \quad (1.42)$$

when $|z_i - z_j|, |r_k - z_j| > |b|$ for $i = 1, \dots, l, i \neq j$ and $k = 1, \dots, n$.

Let $m_{op}^{(n)} := m_{cl-op}^{(0;n)}$. The definition of open-closed field algebra immediately implies that $(V_{op}, m_{op}, \mathbf{d}_{op}, D_{op})$ is a boundary field algebra. In particular, the map Y_{op} defined in (1.5) satisfies the equations (1.6), (1.7), (1.8) and (1.9). Similarly, we know that the map $\mathbb{Y}$ defined in (1.23) satisfies the equations (1.24), (1.25), (1.26), (1.27), (1.28), (1.29), (1.30) and (1.31).

In an open-closed field algebra $(V_{cl}, V_{op}, m_{cl-op})$, there is an additional vertex operator map:

$$\begin{aligned} \mathbb{Y}_{cl-op} : (V_{cl} \otimes V_{op}) \times \mathbb{H} &\rightarrow \overline{V}_{op} \\ (u \otimes v, (z, \bar{z})) &\mapsto \mathbb{Y}_{cl-op}(u; z, \bar{z})v, \end{aligned}$$

defined by

$$\mathbb{Y}_{cl-op}(u; z, \bar{z})v := m_{cl-op}^{(1;1)}(u; v; z, \bar{z}; 0). \quad (1.43)$$

By the convergence property, we have the following identity property:

$$\begin{aligned} \mathbb{Y}_{cl-op}(\mathbf{1}_{cl}; z, \bar{z})v &= m_{cl-op}^{(1;1)}(m_{cl}^{(0)}(1); v; z, \bar{z}; 0) \\ &= m_{cl-op}^{(0;1)}(v; 0) \\ &= v. \end{aligned} \quad (1.44)$$

Proposition 1.11. *For $u \in V_{cl}$, we have*

$$\begin{aligned} [\mathbf{d}_{op}, \mathbb{Y}_{cl-op}(u; z, \bar{z})] &= \mathbb{Y}_{cl-op}((\mathbf{d}_{cl}^L + \mathbf{d}_{cl}^R)u; z, \bar{z}) \\ &\quad + \left(z \frac{\partial}{\partial z} + \bar{z} \frac{\partial}{\partial \bar{z}} \right) \mathbb{Y}_{cl-op}(u; z, \bar{z}). \end{aligned} \quad (1.45)$$

Proof. Applying $\frac{\partial}{\partial a}|_{a=0}$ to both sides of (1.38) when $n = 1, l = 1$ and $r_1 = 0$, we obtain (1.45) immediately. ■

Proposition 1.12. *For $u \in V_{cl}$, we have*

$$[D_{op}, \mathbb{Y}_{cl-op}(u; z, \bar{z})] = \mathbb{Y}_{cl-op}((D_{cl}^L + D_{cl}^R)u; z, \bar{z}), \quad (1.46)$$

$$\mathbb{Y}_{cl-op}(D_{cl}^L u; z, \bar{z}) = \frac{\partial}{\partial z} \mathbb{Y}_{cl-op}(u; z, \bar{z}), \quad (1.47)$$

$$\mathbb{Y}_{cl-op}(D_{cl}^R u; z, \bar{z}) = \frac{\partial}{\partial \bar{z}} \mathbb{Y}_{cl-op}(u; z, \bar{z}). \quad (1.48)$$

Proof. Applying $\frac{\partial}{\partial b}|_{b=0}$ ($\frac{\partial}{\partial \bar{b}}|_{\bar{b}=0}$) to both sides of (1.42) when $n = 1, l = 1$ and $r_1 = 0$, we obtain (1.47) and (1.48).

Applying $\frac{\partial}{\partial a}|_{a=0}$ to both sides of (1.39) when $n = 1, l = 1$ and $r_1 = 0$ and using (1.47) and (1.48) we obtain the first identity in (1.46) immediately. ■

1.4 Analytic open-closed field algebras

The notion of open-closed field algebra introduced in the last subsection is very general. There are not much to say about open-closed field algebras in such generality. In this subsection, we study those open-closed field algebras satisfying some nice analytic properties. In this case, the whole structure can be reconstructed from some simple ingredients.

Definition 1.13. An open-closed field algebra $(V_{cl}, V_{op}, m_{cl-op})$ is called *analytic* if it satisfies the following conditions:

1. $\mathbb{Y}_{cl-op}$ can be extended to a map $V_{cl} \otimes V_{op} \times \mathbb{H} \times \overline{\mathbb{H}} \rightarrow \overline{V}_{op}$ such that for $z \in \mathbb{H}, \zeta \in \overline{\mathbb{H}}$

$$\mathbb{Y}_{cl-op}(u; z, \bar{z}) = \mathbb{Y}_{cl-op}(u; z, \zeta)|_{\zeta=\bar{z}}. \quad (1.49)$$

2. For $n \in \mathbb{N}$, $v_1, \dots, v_{n+1} \in V_{op}, v' \in (V_{op})'$ and $u_1, \dots, u_n \in V_{cl}$, the series

$$\langle v', \mathbb{Y}_{cl-op}(u_1, z_1, \zeta_1) Y_{op}(v_1, r_1) \cdots \mathbb{Y}_{cl-op}(u_n, z_n, \zeta_n) Y_{op}(v_n, r_n) v_{n+1} \rangle$$

is absolutely convergent when $|z_1|, |\zeta_1| > r_1 > \cdots > |z_n|, |\zeta_n| > r_n > 0$ and can be extended to a (possibly multivalued) analytic function on

$$\{(z_1, \zeta_1, r_1, \dots, z_n, \zeta_n, r_n) \in M_{\mathbb{C}}^{3n}\}.$$

3. For $n \in \mathbb{N}$, $v', u_1, \dots, u_{n+1} \in V_{cl}$, the series

$$\langle v', \mathbb{Y}_{an}(u_1; z_1, \zeta_1) \cdots \mathbb{Y}_{an}(u_n; z_n, \zeta_n) u_{n+1} \rangle$$

is absolutely convergent when $|z_1| > \cdots > |z_n| > 0$ and $|\zeta_1| > \cdots > |\zeta_n| > 0$ and can be extended to an analytic function on $M_{\mathbb{C}}^{2n}$.

4. For $v' \in V'_{op}, v_1, v_2 \in V_{op}, u \in V_{cl}$, the series

$$\langle v', Y_{op}(\mathbb{Y}_{cl-op}(u; z, \zeta)v_1, r)v_2 \rangle$$

is absolutely convergent when $r > |z|, |\zeta| > 0$.

5. For $v' \in V'_{op}, v \in V_{op}, u_1, u_2 \in V_{cl}$, the series

$$\langle v', \mathbb{Y}_{cl-op}(\mathbb{Y}_{an}(u_1; z_1, \zeta_1)u_1; z_2, \zeta_2)v \rangle,$$

converges absolutely when $|z_2| > |z_1| > 0, |\zeta_2| > |\zeta_1| > 0$ and $|z_1| + |\zeta_1| < |z_2 - \zeta_2|$.

By the convergence properties of open-closed field algebra, $m_{cl-op}^{(l;n)}$ can be expressed as products of $\mathbb{Y}_{cl-op}, Y_{op}$ on a dense subdomain of $M_{\mathbb{H}}^l \times \Lambda^n$. $m_{cl-op}^{(l;n)}$ on the complement of this dense subdomain is completely determined by analytic extension. Therefore, $m_{cl-op}^{(l;n)}, l, n \in \mathbb{N}$ are completely determined by $\mathbf{1}_{op}, Y_{op}$ and $\mathbb{Y}_{cl-op}$. Similarly, $m_{cl}^{(n)}, n \in \mathbb{N}$ is completely determined by $\mathbf{1}_{cl}$ and $\mathbb{Y}_{an}$. Therefore, we also denote an analytic open-closed conformal field algebra by

$$((V_{cl}, \mathbb{Y}_{an}, \mathbf{1}_{cl}), (V_{op}, Y_{op}, \mathbf{1}_{op}), \mathbb{Y}_{cl-op}).$$

or $(V_{cl}, V_{op}, \mathbb{Y}_{cl-op})$ for simplicity.

Two immediate consequences of the definition of analytic open-closed field algebra are given in the following two Lemmas.

Lemma 1.14. *D_{op} -bracket properties: $u \in V_{cl}$, we have*

$$\begin{aligned} [D_{op}, \mathbb{Y}_{cl-op}(u; z, \zeta)] &= \mathbb{Y}_{cl-op}((D_{cl}^L + D_{cl}^R)u; z, \zeta), \\ \mathbb{Y}_{cl-op}(D_{cl}^L u; z, \zeta) &= \frac{\partial}{\partial z} \mathbb{Y}_{cl-op}(u; z, \zeta), \\ \mathbb{Y}_{cl-op}(D_{cl}^R u; z, \zeta) &= \frac{\partial}{\partial \zeta} \mathbb{Y}_{cl-op}(u; z, \zeta). \end{aligned} \quad (1.50)$$

Proof. By (1.46), (1.47) and (1.48), for any fixed $z \in \mathbb{H}$, all three identities holds when $\zeta = \bar{z}$. Replacing u by $(D_{cl}^R)^k u$ for any $k \in \mathbb{N}$ in (1.46), (1.47) and (1.48), we obtained that all derivatives $(\frac{\partial}{\partial \zeta})^k|_{\zeta=\bar{z}}, \forall k \in \mathbb{N}$ of both sides of above three identities equal respectively. By the property of analytic function and the fact that $\overline{\mathbb{H}}$ is a simply connected domain, we obtain that both sides of above three identities, viewed as analytic functions for a fixed $z \in \mathbb{H}$ and any $\zeta \in \overline{\mathbb{H}}$, must equal respectively. ■

Lemma 1.15. *$\mathbf{d}_{op}$ -bracket properties : $u \in V_{cl}$, we have*

$$\begin{aligned} [\mathbf{d}_{op}, \mathbb{Y}_{cl-op}(u; z, \zeta)] &= \mathbb{Y}_{cl-op}((\mathbf{d}_{cl}^L + \mathbf{d}_{cl}^R)u; z, \zeta) \\ &\quad + \left(z \frac{\partial}{\partial z} + \zeta \frac{\partial}{\partial \zeta} \right) \mathbb{Y}_{cl-op}(u; z, \zeta). \end{aligned} \quad (1.51)$$

Proof. The proof is the same as that of Lemma 1.14. ■

An analytic open-closed field algebra $(V_{cl}, V_{op}, \mathbb{Y}_{cl-op})$ satisfies two associativities and two commutativities. These properties are very important for categorical formulation.

Proposition 1.16 (Associativity I). *For $u \in V_{cl}, v_1, v_2 \in V_{op}, v' \in V'_{op}$ and $z \in \mathbb{H}, \zeta \in \overline{\mathbb{H}}$, we have*

$$\langle v', \mathbb{Y}_{cl-op}(u; z, \zeta) Y_{op}(v_1, r) v_2 \rangle = \langle v', Y_{op}(\mathbb{Y}_{cl-op}(u; z - r, \zeta - r) v_1, r) v_2 \rangle \quad (1.52)$$

when $|z|, |\zeta| > r > 0$ and $r > |r - z|, |r - \zeta| > 0$.

Proof. We abbreviate the “left (right) hand side” as “LHS (RHS)” . For $z \in \mathbb{H}$,

$$\begin{aligned} \text{LHS of (1.52)}|_{\zeta=\bar{z}} &= m_{cl-op}^{(1;1)}(u; m_{op}^{(2)}(v_1, v_2; r, 0); z, \bar{z}; 0) \\ &= m_{cl-op}^{(1;2)}(u; v_1, v_2; z, \bar{z}; r, 0) \end{aligned} \quad (1.53)$$

when $|z| > r > 0$, and

$$\begin{aligned} \text{RHS of (1.52)}|_{\zeta=\bar{z}} &= m_{cl-op}^{(0;2)}(m_{cl-op}^{(1;1)}(u; v_1; z - r, \bar{z} - r; 0), v_2; r, 0) \\ &= m_{cl-op}^{(1;2)}(u; v_1, v_2; z, \bar{z}; r, 0) \end{aligned} \quad (1.54)$$

when $r > |r - z| > 0$. Therefore (1.52) holds when $\zeta = \bar{z}$ and $|z| > r > |r - z| > 0, z \in \mathbb{H}$. Now replace u in (1.52) by $(D_{cl}^R)^k u, k \in \mathbb{N}$, we obtain:

$$\left. \frac{\partial^k}{\partial \zeta^k} \right|_{\zeta=\bar{z}} \text{LHS of (1.52)} = \left. \frac{\partial^k}{\partial \zeta^k} \right|_{\zeta=\bar{z}} \text{RHS of (1.52)} \quad (1.55)$$

when $z \in \mathbb{H}$ and $|z| > r > |r - z| > 0$. Then by the properties of analytic function, it is clear that (1.52) holds for all $z \in \mathbb{H}, \zeta \in \overline{\mathbb{H}}$ and $|z|, |\zeta| > r > 0$ and $r > |r - z|, |r - \zeta| > 0$. $\blacksquare$

Proposition 1.17 (Associativity II). *For $u_1, u_2 \in V_{cl}, v_1, v_2 \in V_{op}, v' \in V'_{op}$ and $z_1, z_2 \in \mathbb{H}, \zeta_1, \zeta_2 \in \overline{\mathbb{H}}$, we have*

$$\begin{aligned} \langle v', \mathbb{Y}_{cl-op}(u_1; z_1, \zeta_1) \mathbb{Y}_{cl-op}(u_2; z_2, \zeta_2) v_2 \rangle \\ = \langle v', \mathbb{Y}_{cl-op}(\mathbb{Y}_{an}(u_1; z_1 - z_2, \zeta_1 - \zeta_2) u_2; z_2, \zeta_2) v_2 \rangle \end{aligned} \quad (1.56)$$

when $|z_1|, |\zeta_1| > |z_2|, |\zeta_2|$ and $|z_2| > |z_1 - z_2| > 0, |\zeta_2| > |\zeta_1 - \zeta_2| > 0$ and $|z_2 - \zeta_2| > |z_1 - z_2| + |\zeta_1 - \zeta_2|$.

Proof. The proof is similar to that of (1.52). So we will only sketch the difference here. First, it is easy to show that (1.56) is true for $z_i \in \mathbb{H}, \zeta_i = \bar{z}_i, i = 1, 2$ in a proper domain. Then replacing u_1, u_2 by $(D_{cl}^R)^p u_1, (D_{cl}^R)^q u_2$ for $p, q \in \mathbb{N}$ and using (1.29) and (1.50), we obtain that

$$\left. \frac{\partial^p}{\partial \zeta_1^p} \frac{\partial^q}{\partial \zeta_2^q} \right|_{\zeta_i=\bar{z}_i} \text{LHS of (1.56)} = \left. \frac{\partial^p}{\partial \zeta_1^p} \frac{\partial^q}{\partial \zeta_2^q} \right|_{\zeta_i=\bar{z}_i} \text{RHS of (1.56)} \quad (1.57)$$

in a proper domain. By the property of analytic function again, the analytic extension of both sides of (1.56) to the following simply connected domain

$$D_0 = \{(z_1, \zeta_1, z_2, \zeta_2) | z_i \in \mathbb{H}, \zeta_i \in \overline{\mathbb{H}}, |z_1|, |\zeta_1| > |z_2|, |\zeta_2|\}$$

must be identical. Moreover, the additional restrictions of domain in the statement of Associativity II guarantee the absolute convergence of both sides of (1.56). $\blacksquare$

For an analytic open-closed field algebra, the map $\mathbb{Y}_{cl-op}$ can be uniquely extended to $V_{cl} \otimes V_{op} \times R$ where

$$R := \{(z, \zeta) \in \mathbb{C}^2 | z \in \mathbb{H} \cup \mathbb{R}_+, \zeta \in \overline{\mathbb{H}} \cup \mathbb{R}_+, z \neq \zeta\}.$$


Proposition 1.18 (Commutativity I). For $u \in V_{cl}, v_1, v_2 \in V_{op}$ and $v' \in V'_{op}$,

$$\langle v', \mathbb{Y}_{cl-op}(u; z, \zeta) Y_{op}(v_1, r) v_2 \rangle, \quad (1.58)$$

which is absolutely convergent when $z > \zeta > r > 0$ (recall Definition 1.13), and

$$\langle v', Y_{op}(v_1, r) \mathbb{Y}_{cl-op}(u; z, \zeta) v_2 \rangle, \quad (1.59)$$

which is absolutely convergent when $r > z > \zeta > 0$ (recall Definition 1.13), are analytic continuation of each other along the following path ².



Proof. When $z \in \mathbb{H}$, $\zeta = \bar{z}$ and $|z| > r > 0$, (1.58) is absolutely convergent to $m_{cl-op}^{(1;2)}(u; v_1, v_2; z, \bar{z}; r, 0)$. Hence, if we analytic extend the analytic function (1.58) to

$$D_1 := \{(z, \zeta) | z \in \mathbb{H}, \zeta = \bar{z}, |z| = r\},$$

then its value on D_1 must equal to $m_{cl-op}^{(1;2)}(u; v_1, v_2; z, \bar{z}; r, 0)$ when $z \in \mathbb{H}, |z| = r$ by the continuity of $m_{cl-op}^{(1;2)}$.

Similarly, the unique extension of (1.59) from $r > |z|, |\zeta| > 0$ to D_1 also equal to $m_{cl-op}^{(1;2)}(u; v_1, v_2; z, \bar{z}; r, 0)$ when $z \in \mathbb{H}, |z| = r$.

Both (1.58) and (1.59) can be uniquely extended to two analytic functions on $\mathbb{H} \times \overline{\mathbb{H}}$. By the discussion above, these two extended analytic functions of (z, ζ) take the same value on D_1 which is of lower dimension.

Now we replace u by $(D_{cl}^L)^m (D_{cl}^R)^n u, m, n \in \mathbb{N}$ and repeat the arguments in the proof of Proposition 1.16. We obtain that all the derivatives of the two extended analytic functions also match on D_1 . By the properties of analytic functions, these two extended functions are identical on $\mathbb{H} \times \overline{\mathbb{H}}$. Then their extensions to $(z, \zeta) \in R \setminus \{z = r, \text{ or } \zeta = r\}$ are also identical. Thus we have proved the first commutativity. ■


Proposition 1.19 (Commutativity II). For $u_1, u_2 \in V_{cl}, v \in V_{op}$ and $v' \in V'_{op}$,

$$\langle v', \mathbb{Y}_{cl-op}(u_1; z_1, \zeta_1) \mathbb{Y}_{cl-op}(u_2; z_2, \zeta_2) v \rangle, \quad (1.61)$$

which is absolutely convergent when $z_1 > \zeta_1 > z_2 > \zeta_2 > 0$, and

$$\langle v', \mathbb{Y}_{cl-op}(u_2; z_2, \zeta_2) \mathbb{Y}_{cl-op}(u_1; z_1, \zeta_1) v \rangle, \quad (1.62)$$

which is absolutely convergent when $z_2 > \zeta_2 > z_1 > \zeta_1 > 0$, are analytic continuation of each other along the following paths.



²The extended domain $R \setminus \{z = r, \text{ or } \zeta = r\}$ is simply connected for fixed $r > 0$, all possible paths of analytic continuation are homotopically equivalent.

Proof. Commutativity II follows directly from the Associativity II (1.56) and the skew-symmetry of the full field algebra V_{cl} [HKo2]. Here, we give a more direct proof which is similar to that of Proposition 1.18. The unique extension of (1.61) from $|z_1|, |\zeta_1| > |z_2|, |\zeta_2| > 0$ to

$$D_2 := \{(z_1, \zeta_1, z_2, \zeta_2) | z_1, z_2 \in M_{\mathbb{H}}^2, |z_1| = |z_2|, \zeta_i = \bar{z}_i, i = 1, 2\}$$

equal to $m_{cl-op}^{(2;1)}(u_1, u_2; v; z_1, \bar{z}_1, z_2, \bar{z}_2; 0)$, the unique extension of (1.62) from $|z_2|, |\zeta_2| > |z_1|, |\zeta_1| > 0$ to D_2 also match with $m_{cl-op}^{(2;1)}(u_1, u_2; v; z_1, \bar{z}_1, z_2, \bar{z}_2; 0)$. By the similar argument as the proof of the first commutativity, we see that (1.61) in $|z_1|, |\zeta_1| > |z_2|, |\zeta_2| > 0$ and (1.61) in $|z_2|, |\zeta_2| > |z_1|, |\zeta_1| > 0$ are analytic continuation of each other along the following paths.

(1.64)

Then it is obvious to see that the unique extension of both (1.61) and (1.62) to the subdomain of $\{(z_1, \zeta_1, z_2, \zeta_2) \in \mathbb{R}_+^4\}$, where $z_1, \zeta_1, z_2, \zeta_2$ have distinct values, are analytic extension of each other along the paths (1.63). ■

1.5 Open-closed field algebras over U

We gradually add more structures on an analytic open-closed field algebra. At the end of this process, we will arrive at a notion called open-closed field algebra over a vertex operator algebra U . Let $(V^L, Y_{V^L}, \mathbf{1}^L)$ and $(V^R, Y_{V^R}, \mathbf{1}^R)$ be two vertex algebras. We now consider an analytic open-closed field algebra $(V_{cl}, V_{op}, \mathbb{Y}_{cl-op})$ such that V_{cl} is a full field algebra over $V^L \otimes V^R$. For such full field algebra V_{cl} , we will not distinguish V^L with $V^L \otimes \mathbf{1}^R \subset V_{cl}$ and V^R with $\mathbf{1}^L \otimes V^R \subset V_{cl}$.

Lemma 1.20. *For $u^L \in V^L$ and $u^R \in V^R$, $\mathbb{Y}_{cl-op}(u^L; z, \zeta)$ is independent of ζ and $\mathbb{Y}_{cl-op}(u^R; z, \zeta)$ is independent of z .*

Proof. For $u^L \in V^L, w \in V_{op}, w' \in (V_{op})'$, using the associativity (1.56), we have

$$\begin{aligned} \langle w', \mathbb{Y}_{cl-op}(u^L; z, \zeta)w \rangle &= \langle w', \mathbb{Y}_{cl-op}(u^L; z, \zeta) \mathbb{Y}_{cl-op}(\mathbf{1}_{cl}; z_1, \zeta_1)w \rangle \\ &= \langle w', \mathbb{Y}_{cl-op}(\mathbb{Y}_{an}(u^L; z - z_1, \zeta - \zeta_1) \mathbf{1}_{cl}; z_1, \zeta_1)w \rangle \end{aligned} \quad (1.65)$$

when $|z_1| > |z - z_1| > 0$, $|\zeta_1| > |\zeta - \zeta_1| > 0$ and $|z - z_1| + |\zeta - \zeta_1| < |z_1 - \zeta_1|$. The right hand side of (1.65) is independent of ζ and the left hand side of (1.65) is analytic in ζ . Hence $\mathbb{Y}_{cl-op}(u^L; z, \zeta)$ is independent of ζ for all $z \in \mathbb{H}$. Similarly, $\mathbb{Y}_{cl-op}(u^R; z, \zeta)$ is independent of z for all $\zeta \in \mathbb{H}$ and $u^R \in V^R$. ■

In order to emphasis these ζ - or z -independence properties, we denote them simply as $\mathbb{Y}_{cl-op}(u^L, z)$ and $\mathbb{Y}_{cl-op}(u^R, \zeta)$ for $u^L \in V^L$ and $u^R \in V^R$ respectively.

Replace u in (1.51) by $u^L \in V^L$ and $u^R \in V^R$ respectively, we obtain

$$\begin{aligned} [\mathbf{d}_{op}, \mathbb{Y}_{cl-op}(u^L, z)] &= \mathbb{Y}_{cl-op}(\mathbf{d}_{cl}^L u^L, z) + z \frac{\partial}{\partial z} \mathbb{Y}_{cl-op}(u^L, z), \\ [\mathbf{d}_{op}, \mathbb{Y}_{cl-op}(u^R, \zeta)] &= \mathbb{Y}_{cl-op}(\mathbf{d}_{cl}^R u^R, \zeta) + \zeta \frac{\partial}{\partial \zeta} \mathbb{Y}_{cl-op}(u^R, \zeta). \end{aligned} \quad (1.66)$$

As a consequence, we have

$$\begin{aligned}\mathbb{Y}_{cl-op}(u^L, z) &= \sum_{n \in \mathbb{R}} u_n^L z^{-n-1}, \\ \mathbb{Y}_{cl-op}(u^R, \zeta) &= \sum_{n \in \mathbb{R}} u_n^R \zeta^{-n-1},\end{aligned}\tag{1.67}$$

where $u_n^L, u_n^R \in \text{End } V_{op}$ and $\text{wt } u_n^L = \text{wt } u^L - n - 1$ and $\text{wt } u_n^R = \text{wt } u^R - n - 1$, and $z^n = e^{n \log z}$ and $\zeta^n = e^{n \log \zeta}$.

Moreover, we have

$$\begin{aligned}[D_{op}, \mathbb{Y}_{cl-op}(u^L, z)] &= \mathbb{Y}_{cl-op}(D_{cl}^L u^L, z) = \frac{\partial}{\partial z} \mathbb{Y}_{cl-op}(u^L, z), \\ [D_{op}, \mathbb{Y}_{cl-op}(u^R, \zeta)] &= \mathbb{Y}_{cl-op}(D_{cl}^R u^R, \zeta) = \frac{\partial}{\partial \zeta} \mathbb{Y}_{cl-op}(u^R, \zeta),\end{aligned}\tag{1.68}$$

which further implies that

$$\begin{aligned}e^{aD_{op}} \mathbb{Y}_{cl-op}(u^L, z) e^{-aD_{op}} &= \mathbb{Y}_{cl-op}(u^L, z + a) \\ e^{aD_{op}} \mathbb{Y}_{cl-op}(u^R, \zeta) e^{-aD_{op}} &= \mathbb{Y}_{cl-op}(u^R, \zeta + a)\end{aligned}\tag{1.69}$$

for $|z| > |a|, z + a \in \mathbb{H}$ and $|\zeta| > |a|, \zeta + a \in \overline{\mathbb{H}}$ respectively.

Lemma 1.21. *For any $u^L \in V^L \otimes \mathbf{1}^R$ and $u^R \in \mathbf{1}^L \otimes V^R$, the following two limits:*

$$\lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op}, \quad \lim_{\zeta \rightarrow 0} \mathbb{Y}_{cl-op}(u^R, \zeta) \mathbf{1}_{op}$$

exist in V_{op} .

Proof. By the associativity and the creation property of open-string vertex algebra, we have

$$\begin{aligned}\mathbb{Y}_{cl-op}(u^L, z + r) \mathbf{1}_{op} &= \mathbb{Y}_{cl-op}(u^L, z + r) Y_{op}(\mathbf{1}_{op}, r) \mathbf{1}_{op} \\ &= Y_{op}(\mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op}, r) \mathbf{1}_{op} \\ &= e^{rD_{op}} \mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op}\end{aligned}\tag{1.70}$$

when $|z + r| > r > |z| > 0$. For fixed $z \in \mathbb{H}$, the left hand side of (1.70) is an analytic function valued in $\overline{V}_{op}$ on the domain $\{r \in \mathbb{C} | z + r \neq 0\}$. The right hand side of (1.70), as a power series of r , is absolutely convergent when $|r| > |z| > 0$. By the general property of power series, the right hand side of (1.70) is absolutely convergent for all $r \in \mathbb{C}$ to a singlevalued analytic function. Because both sides of (1.70) are analytic functions, the equality (1.70) must hold for all $r \in \mathbb{C}$. In particular, $\lim_{r \rightarrow -z} \mathbb{Y}_{cl-op}(u^L, z + r) \mathbf{1}_{op}$ exists. Equivalently, $\lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op}$ exists. By the expansion (1.67), we must have $u_n^L \mathbf{1}_{op} = 0$ for all $n > -1$. Moreover, it is also easy to see that $u_n^L \mathbf{1}_{op} = 0$ for any $n \notin -\mathbb{Z}_+$ by (1.68) (see the proof of Proposition 1.8 in [HKo1]). Therefore, we have

$$\lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op} = u_{-1}^L \mathbf{1}_{op} \in V_{op}.$$

The proof of the existence of the second limit is entirely the same. ■

By the Lemma above, we can define two maps $h^L : V^L \rightarrow V_{op}$ and $h^R : V^R \rightarrow V_{op}$ as follows: for all $u^L \in V^L$ and $u^R \in V^R$,

$$\begin{aligned} h^L : u^L &\mapsto \lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op}, \\ h^R : u^R &\mapsto \lim_{\zeta \rightarrow 0} \mathbb{Y}_{cl-op}(u^R, \zeta) \mathbf{1}_{op}. \end{aligned} \quad (1.71)$$

Notice also that h^L, h^R preserve the weights. Namely

$$\begin{aligned} \text{wt } h^L(u^L) &= \text{wt } (u^L_{-1} \mathbf{1}_{op}) = \text{wt}^L u^L, \\ \text{wt } h^R(u^R) &= \text{wt } (u^R_{-1} \mathbf{1}_{op}) = \text{wt}^R u^R. \end{aligned}$$

Therefore both h^L and h^R can be naturally extended to maps $\overline{V^L} \rightarrow \overline{V_{op}}$ and $\overline{V^R} \rightarrow \overline{V_{op}}$. We still denote the extended maps as h^L and h^R respectively.

Lemma 1.22. *For $u^L \in V^L, u^R \in V^R$, we have*

$$\begin{aligned} \mathbb{Y}_{cl-op}(u^L, z) \mathbf{1}_{op} &= e^{zD_{op}} h^L(u^L), \\ \mathbb{Y}_{cl-op}(u^R, \zeta) \mathbf{1}_{op} &= e^{\zeta D_{op}} h^R(u^R). \end{aligned} \quad (1.72)$$

Proof. Since we have shown that (1.70) holds for all $r \in \mathbb{C}, z \in \mathbb{H}$ and both sides of (1.70) are analytic for all $r, z \in \mathbb{C}$, if we take the limit $\lim_{z \rightarrow 0}$ on both sides of (1.70), the equality should still hold. Thus we obtain the first identity in (1.72).

The proof of the second identity is entirely the same. ■

Proposition 1.23. *h^L and h^R are homomorphisms from V^L and V^R respectively to their images, viewed as graded vertex algebras.*

Proof. We have shown that h^L, h^R preserve gradings. By the identity property of open-closed field algebra, we have

$$h^L(\mathbf{1}_{cl}) = \mathbf{1}_{op}.$$

Next, for $u^L \in V^L$ and $r > 0$, we define

$$\begin{aligned} \mathbb{Y}_{cl-op}(u^L, r)w &:= \lim_{z \rightarrow r} \mathbb{Y}_{cl-op}(u^L, z)w \\ &= \lim_{z \rightarrow r} \mathbb{Y}_{cl-op}(u^L, z)Y_{op}(\mathbf{1}_{op}, r)w \end{aligned} \quad (1.73)$$

where the limit is taken along a path from a fixed initial point in $\mathbb{H}$ to $r > 0$. Since $\mathbb{Y}_{cl-op}(u^L, z)w$ is analytic in $\mathbb{C}^\times$, the limit is independent of the path we choose. So we choose a path in the domain $\{z \in \mathbb{H} \mid |z| > r > |z - r| > 0\}$. In this domain, we can apply the associativity (1.52) to the right hand side of (1.73). We obtain

$$\begin{aligned} \mathbb{Y}_{cl-op}(u^L, r)w &= \lim_{z \rightarrow r} Y_{op}(\mathbb{Y}_{cl-op}(u^L, z - r) \mathbf{1}_{op}, r)w \\ &= Y_{op}(h^L(u^L), r)w. \end{aligned}$$

For $|z| > r > 0$, by (1.72), we have,

$$\mathbb{Y}_{cl-op}(\mathbb{Y}(u^L; r, r)v^L, z) \mathbf{1}_{op} = e^{zD_{op}} h^L(\mathbb{Y}(u^L; r, r)v^L), \quad (1.74)$$

the right hand side of which is absolutely convergent for all $z \in \mathbb{C}$. And both sides are analytic in z . Therefore $\mathbb{Y}_{cl-op}(\mathbb{Y}(u^L; r, r)v^L, z)\mathbf{1}_{op}$ is absolutely convergent for all $z \in \mathbb{C}$ and (1.74) holds for all $z \in \mathbb{C}$. By the associativity, we have

$$\mathbb{Y}_{cl-op}(\mathbb{Y}(u^L; r, r)v^L, z)\mathbf{1}_{op} = \mathbb{Y}_{cl-op}(u^L, r+z)\mathbb{Y}_{cl-op}(v^L, z)\mathbf{1}_{op} \quad (1.75)$$

for $|r+z| > |z| > r > 0$. Again both sides of (1.75) are analytic in z . Hence the left hand side of (1.75) defined for all $z \in \mathbb{C}$ is the analytic extension of the right hand side of (1.75), which is defined on $\{|r+z| > |z|\}$. Since the extension is free of singularity on entire $\mathbb{C}$, the right hand side of (1.75) must be well-defined on entire $\mathbb{C}$. Therefore, we must have

$$\lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(\mathbb{Y}(u^L, r)v^L, z)\mathbf{1}_{op} = \lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u^L, r+z)\mathbb{Y}_{cl-op}(v^L, z)\mathbf{1}_{op}. \quad (1.76)$$

Combining above results, we have

$$\begin{aligned} h^L(\mathbb{Y}(u^L, r)v^L) &= \lim_{z \rightarrow 0} e^{zD_{op}} h^L(\mathbb{Y}(u^L, r)v^L) \\ &= \lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(\mathbb{Y}(u^L, r)v^L, z)\mathbf{1}_{op} \\ &= \lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u^L, r+z)\mathbb{Y}_{cl-op}(v^L, z)\mathbf{1}_{op} \\ &= \mathbb{Y}_{cl-op}(u^L, r)h^L(v^L) \\ &= Y_{op}(h^L(u^L), r)h^L(v^L). \end{aligned}$$

Thus h^L is a vertex algebra homomorphism. The proof for h^R is entirely the same. $\blacksquare$

Let $(U, Y, \mathbf{1}, \omega)$ be a vertex operator algebra with central charge c . U and $U \otimes U$ naturally give an analytic open-closed field algebra, in which $\mathbb{Y}_{cl-op}(\cdot; z, \zeta) \cdot$ is given by

$$\begin{aligned} \mathbb{Y}_{cl-op}(u \otimes v; z, \zeta)w &= Y(u, z)Y(v, \zeta)w, & |z| > |\zeta| > 0, \\ &= Y(v, \zeta)Y(u, z)w, & |\zeta| > |z| > 0 \end{aligned} \quad (1.77)$$

for $u, v, w \in U$. In this case, $h^L : u \otimes \mathbf{1} \mapsto u$ and $h^R : \mathbf{1} \otimes u \mapsto u$. We denote this open-closed field algebra as $(U \otimes U, U)$.

In general, let $\rho^L, \rho^R \in \text{Aut}(U)$ where $\text{Aut}(U)$ is the set of automorphisms of U as vertex operator algebra. We can obtain a new action of $U \otimes U$ on U by composing (1.77) with the automorphism $\rho^L \otimes \rho^R : U \otimes U \rightarrow U \otimes U$. Namely, there exists another open-closed field algebra structure on U and $U \otimes U$, in which $\mathbb{Y}_{cl-op}(u \otimes v; z, \zeta)w$, for $u, v, w \in U$, is given by

$$\begin{aligned} Y(\rho^L(u), z)Y(\rho^R(v), \zeta)w, & \quad \text{for } |z| > |\zeta| > 0, \\ Y(\rho^R(v), \zeta)Y(\rho^L(u), z)w, & \quad \text{for } |\zeta| > |z| > 0. \end{aligned} \quad (1.78)$$

In this case, $h^L : u \otimes \mathbf{1} \mapsto \rho^L(u)$ and $h^R : \mathbf{1} \otimes u \mapsto \rho^R(u)$. We denote such open-closed field algebra as $(U \otimes U, U, \rho^L, \rho^R)$. In particular, $(U \otimes U, U, \text{id}_U, \text{id}_U)$ is just $(U \otimes U, U)$.

Remark 1.24. $(U \otimes U, U, \rho^L, \rho^R)$ for general automorphisms ρ^L and ρ^R is very interesting in physics. But it adds some technical subtleties in later formulations. So we postpone its study to future publications. In this work, we focus on $(U \otimes U, U)$.

Definition 1.25. Let $(U, Y, \mathbf{1}, \omega)$ be a vertex operator algebra. An *open-closed field algebra over U* is an analytic open-closed field algebra

$$((V_{cl}, m_{cl}, \iota_{cl}), (V_{op}, Y_{op}, \iota_{op}), \mathbb{Y}_{cl-op}),$$

where $(V_{cl}, m_{cl}, \iota_{cl})$ is a conformal full field algebra over $U \otimes U$ and $(V_{op}, Y_{op}, \iota_{op})$ is an open-string vertex operator algebra over U , satisfying the following conditions:

1. *U -invariant boundary condition:* $h^L = h^R = \iota_{op}$.
2. *Chirality splitting property:* $\forall u \in V_{cl}$, $u = u^L \otimes u^R \in W^L \otimes W^R \subset V_{cl}$ for some U -modules W^L, W^R . There exist U -modules W_1, W_2 and intertwining operators $\mathcal{Y}^{(1)}, \mathcal{Y}^{(2)}, \mathcal{Y}^{(3)}, \mathcal{Y}^{(4)}$ of type $\binom{V_{op}}{W^L W_1}, \binom{W_1}{W^R V_{op}}, \binom{V_{op}}{W^R W_2}, \binom{W_2}{W^L V_{op}}$ respectively³, such that

$$\langle w', \mathbb{Y}_{cl-op}(u; z, \zeta)w \rangle = \langle w', \mathcal{Y}^{(1)}(u^L, z) \mathcal{Y}^{(2)}(u^R, \zeta)w \rangle \quad (1.79)$$

when $|z| > |\zeta| > 0$ (recall the convention (1.16) and (1.17)), and

$$\langle w', \mathbb{Y}_{cl-op}(u; z, \zeta)w \rangle = \langle w', \mathcal{Y}^{(3)}(u^R, \zeta) \mathcal{Y}^{(4)}(u^L, z)w \rangle \quad (1.80)$$

when $|\zeta| > |z| > 0$ for all $u \in V_{cl}, w \in V_{op}, w' \in V_{op}$.

In the case that U is generated by ω , i.e. $U = \langle \omega \rangle$, the $\langle \omega \rangle$ -invariant boundary condition is simply called *conformal invariant boundary condition*. We also call open-closed field algebra over $\langle \omega \rangle$ *open-closed conformal field algebra*.

Remark 1.26. The U -invariant boundary condition actually says that the open-closed field algebra over U contains $(U \otimes U, U)$ as a subalgebra. If we only want to construct algebras over Swiss-cheese partial operad, the U -invariant boundary condition in Definition 1.25 can be weakened to the *conformal invariant boundary condition*:

$$h^L|_{\langle \omega \rangle} = h^R|_{\langle \omega \rangle} = \iota_{op}|_{\langle \omega \rangle}. \quad (1.81)$$

These situations appear in physics in the study of the so-called symmetry breaking boundary conditions (see for example [FS1][FS2] and references therein). All examples studied in this work and [K3] satisfy the U -invariant boundary condition. We leave the study of general symmetry-broken situations to the future.

Remark 1.27. The chirality splitting property is a very natural condition because the interior sewing operation of $\tilde{\mathfrak{S}}^c$ is defined by a double sewing operation as given in (2.25). Unfortunately, we do not know whether this chirality splitting property is necessary for general constructions of algebras over $\tilde{\mathfrak{S}}^c$.

For an open-closed field algebra over V , there are three Virasoro elements, $\omega_{op} := \iota_{op}(\omega)$ and $\omega^L := \iota_{cl}(\omega \otimes \mathbf{1})$ and $\omega^R := \iota_{cl}(\mathbf{1} \otimes \omega)$, and we have

$$\begin{aligned} Y_{op}(\omega_{op}, r) &= \sum_{n \in \mathbb{Z}} L(n) r^{-n-2}, \\ \mathbb{Y}(\omega^L, z) &= \sum_{n \in \mathbb{Z}} L^L(n) z^{-n-2}, \\ \mathbb{Y}(\omega^R, \zeta) &= \sum_{n \in \mathbb{Z}} L^R(n) \zeta^{-n-2}. \end{aligned}$$

³It was proved in [HKo1] that V_{op} is a U -modules, and in [HKo2] that V_{cl} is a $U \otimes U$ -modules.

where $L^L(n) = L(n) \otimes 1$ and $L^R(n) = 1 \otimes L(n)$ for $n \in \mathbb{Z}$.

When $U = V$ a vertex operator algebra satisfying the conditions in Theorem 0.1, we have a very simple description of open-closed field algebra over V given in the following Theorem.

Theorem 1.28. *An open-closed field algebra over V is equivalent to the following structure: $(V_{op}, Y_{op}, \iota_{op})$ an open-string vertex operator algebra over V and $(V_{cl}, m_{cl}, \iota_{cl})$ a conformal full field algebra over $V \otimes V$, together with a vertex map $\mathbb{Y}_{cl-op}(\cdot; z, \zeta)$ given by intertwining operators $\mathcal{Y}^{(i)}, i = 1, 2, 3, 4$ as in (1.79) and (1.80), satisfying the unit property:*

$$\mathbb{Y}_{cl-op}(\mathbf{1}_{cl}; z, \bar{z})v := \mathcal{Y}^{(1)}(\mathbf{1}, z)\mathcal{Y}^{(2)}(\mathbf{1}, \bar{z})v = v, \quad \forall v \in V_{op}, \quad (1.82)$$

Associativity I (1.52), Associativity II (1.56) and Commutativity I given in Proposition 1.18.

Proof. It is clear that an open-closed field algebra over V gives the data and properties included in the statement of the Theorem. We only need to show that such data is sufficient to reconstruct an open-closed field algebra over V . Moreover, such open-closed field algebra over V with the given data is unique.

Since V satisfies the condition in Theorem 0.1, the conditions listed in the Definition 1.13 are all automatically satisfied. In particular, for $v_i \in V_{op}, u_i \in V_{cl}, i = 1, \dots, n$ and $v' \in V'_{op}$, the following series:

$$\langle v', \mathbb{Y}_{cl-op}(u_1; z_1, \zeta_1)Y_{op}(v_1, r_1) \cdots \mathbb{Y}_{cl-op}(u_n; z_n, \zeta_n)Y_{op}(v_n, r_n)\mathbf{1}_{op} \rangle. \quad (1.83)$$

is absolutely convergent when $z_1 > \zeta_1 > r_1 > \cdots > z_n > \zeta_n > r_n > 0$, and can be extended by analytic continuation to a multi-valued analytic function for variables $z_i, \zeta_i \in \mathbb{C}^\times, i = 1, \dots, n$ with possible singularities only when two of z_i, ζ_j, r_k are equal. Using this property of (1.83), we can define a non-analytic but single-valued smooth function in $\Lambda^n \times M_{\mathbb{H}}^n$ as follows.

Let $(r_1, \dots, r_n) \in \Lambda^n$ and $(\xi_1, \dots, \xi_n) \in M_{\mathbb{H}}^n, n \in \mathbb{Z}_+$. Let γ_1 be a smooth path from $(3n, 3(n-1), \dots, 1)$ to $(\xi_1, \dots, \xi_n)$ such that $\gamma_1((0, 1)) \subset M_{\mathbb{H}}^n$. Let $\gamma_0 : [0, 1] \rightarrow \mathbb{R}_+^n$ be so that

$$\gamma_0(t) = ((1-t)(3n-1) + t3n, \dots, (1-t)2 + t3).$$

Clearly, γ_0 is the straight line from $(3n-1, 3(n-1)-1, \dots, 2)$ to $(3n, 3(n-1), \dots, 3)$ in $\mathbb{R}_+^n$. Then we define a path $\gamma_2 : [0, 1] \in (\overline{\mathbb{H}} \cup \mathbb{R}_+)^n$ to be the composition $\gamma_2 = \bar{\gamma}_1 \circ \gamma_0$ where $\bar{\gamma}_1$ is the complex conjugation of γ_1 . It is clear that γ_2 is a path from $(3n-1, 3(n-1)-1, \dots, 2)$ to $(\bar{\xi}_1, \dots, \bar{\xi}_n)$. Let $\gamma_3 : [0, 1] \rightarrow \mathbb{R}_+^n$ be so that

$$\gamma_3(t) = ((1-t)(3n-2) + t3n-1, \dots, (1-t)1 + t3n-2).$$

So γ_3 is the straight line from $(3n-2, 3(n-1)-2, \dots, 1)$ to $(r_1, \dots, r_n) \in \Lambda^n$.

Combining $\gamma_1, \gamma_2, \gamma_3$, we obtain a path γ in $\mathbb{C}^{3n}$ from the initial point $(3n, 3n-1, 3n-2, \dots, 3, 2, 1) \in \mathbb{R}_+^{2n}$ to the final point $(\xi_1, \bar{\xi}_1, r_1, \dots, \xi_n, \bar{\xi}_n, r_n)$ in the obvious way.

Then we define

$$\langle v', m_{cl-op}^{(n;n)}(u_1, \dots, u_n, v_1, \dots, v_n; \xi_1, \bar{\xi}_1, \dots, \xi_n, \bar{\xi}_n, r_1, \dots, r_n) \rangle$$

to be the value obtained from the value of (1.83) at the initial point $(3n, 3n-1, 3n-2, \dots, 3, 2, 1)$ by analytic continuation along the path γ . Following a similar argument as in the proof of Theorem 2.11 in [HKo2], it is easy to show, that such defined

$$m_{cl-op}^{(n;n)}(u_1, \dots, u_n; v_1, \dots, v_n; \xi_1, \bar{\xi}_1, \dots, \xi_n, \bar{\xi}_n, r_1, \dots, r_n), \quad (1.84)$$

is independent of the choice of γ_1 and its initial points. Moreover, such defined (1.84) is single-valued and smooth in $\mathbb{H} \times \overline{\mathbb{H}}$.

For $n > l \geq 0$, we define

$$\begin{aligned} m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_n; \xi_1, \bar{\xi}_1, \dots, \xi_l, \bar{\xi}_l; r_1, \dots, r_n), \\ := m_{cl-op}^{(n;n)}(u_1, \dots, u_l, \mathbf{1}_{cl}, \dots, \mathbf{1}_{cl}; v_1, \dots, v_n; \xi_1, \bar{\xi}_1, \dots, \xi_n, \bar{\xi}_n; r_1, \dots, r_n), \end{aligned}$$

and for $l > n \geq 0$, we define

$$\begin{aligned} m_{cl-op}^{(l;n)}(v_1, \dots, v_l; w_1, \dots, w_n; \xi_1, \bar{\xi}_1, \dots, \xi_l, \bar{\xi}_l; r_1, \dots, r_n), \\ := m_{cl-op}^{(l;l)}(u_1, \dots, u_l; v_1, \dots, v_n, \mathbf{1}_{op}, \dots, \mathbf{1}_{op}; \xi_1, \bar{\xi}_1, \dots, \xi_l, \bar{\xi}_l; r_1, \dots, r_l), \end{aligned} \quad (1.85)$$

and for $n = l = 0$, we define $m_{cl-op}^{(0;0)}(\mathbf{1}) = \mathbf{1}_{op} \in V_{op}$.

Immediately following from the construction of $m_{cl-op}^{(l;n)}$, we have, for all $v \in V_{op}$,

$$m_{cl-op}^{(0;1)}(v; 0) = m_{cl-op}^{(1;1)}(\mathbf{1}_{cl}; v; z; 0) = \mathbb{Y}_{cl-op}(\mathbf{1}_{cl}; z, \bar{z})v = v.$$

Now we show the permutation axiom for $m_{cl-op}^{(l;n)}$. This is enough to just consider adjacent permutations $(ii+1), i = 1, \dots, l-1$ because they generate the whole permutation group. We can just consider (12) because all the other cases are exactly the same. This is amount to show that

$$\begin{aligned} m_{cl-op}^{(l;n)}(v_1, v_2, \dots, v_l; w_1, \dots, w_n; \xi_1, \bar{\xi}_1, \xi_2, \bar{\xi}_2, \dots, \xi_l, \bar{\xi}_l; r_1, \dots, r_n), \\ = m_{cl-op}^{(l;n)}(v_2, v_1, \dots, v_l; w_1, \dots, w_n; \xi_2, \bar{\xi}_2, \xi_1, \bar{\xi}_1, \dots, \xi_l, \bar{\xi}_l; r_1, \dots, r_n). \end{aligned} \quad (1.86)$$

By our construction, the only difference of the two sides of (1.86) is that they are obtained by analytic continuation along paths with different initial points. The initial points of the path for the left hand side of (1.86) is $z_1 > \zeta_1 > r_1 > z_2 > \zeta_2 > r_2 > \dots$, that for the right hand side of (1.86) is $z_2 > \zeta_2 > r_1 > z_1 > \zeta_1 > r_2 > \dots$. But by the commutativity, the value of (1.83) at these two initial points are analytic continuation of each other along the paths given in the commutativity axiom of open-closed field algebra. Hence the equality (1.86) follows.

((1.38) and (1.39)) can be proved by first proving similar properties of (1.83), which is obvious by the properties of intertwining operators. Then those properties (1.38) and (1.39) of $m_{cl-op}^{(l;n)}$ follow from analytic continuations.

Since $\mathcal{Y}^{(i)}, i = 1, 2, 3, 4$ are intertwining operators of V , we have, for $u \in V$,

$$\begin{aligned} h^L(u) &= \lim_{z \rightarrow 0} \mathbb{Y}_{cl-op}(u, z) \mathbf{1}_{op} \\ &= \lim_{z \rightarrow 0} \lim_{z_1 \rightarrow 0} \mathcal{Y}^{(1)}(Y(u, z_1) \mathbf{1}, z) \mathcal{Y}^{(2)}(\mathbf{1}, \bar{z}) \iota_{op}(\mathbf{1}) \\ &= \lim_{z \rightarrow 0} \lim_{z_1 \rightarrow 0} Y_{V_{op}}(u, z + z_1) \mathcal{Y}^{(1)}(\mathbf{1}, z) \mathcal{Y}^{(2)}(\mathbf{1}, \bar{z}) \iota_{op}(\mathbf{1}) \\ &= \lim_{z \rightarrow 0} \lim_{z_1 \rightarrow 0} Y_{V_{op}}(u, z + z_1) \mathbb{Y}_{cl-op}(\mathbf{1}_{cl}; z, \bar{z}) \iota_{op}(\mathbf{1}) \\ &= \lim_{z \rightarrow 0} \lim_{z_1 \rightarrow 0} Y_{V_{op}}(u, z + z_1) \iota_{op}(\mathbf{1}) \\ &= \lim_{z \rightarrow 0} \lim_{z_1 \rightarrow 0} \iota_{op}(Y(u, z + z_1) \mathbf{1}) \\ &= \iota_{op}(u). \end{aligned} \quad (1.87)$$

Similarly, one can show that $h^R(u) = \iota_{op}(u), \forall u \in V$. Thus we have proved the V -invariant boundary condition $h^L = h^R = \iota_{op}$.

It remains to show the convergence properties of open-closed field algebra. For the first convergence property ((1.33) and (1.34)), one first consider cases when $z_j^{(i)}, r_p^{(i)}, j = 1, \dots, k, p = 1, \dots, m$ in (1.33) have distinct absolute values and $z_j, r_p, j = 1, \dots, l, p = 1, \dots, n$ in (1.33) have distinct absolute values. In these cases, one can express $m_{cl-op}^{(l;n)}$ and $m_{cl-op}^{(k;m)}$ as products of $\mathbb{Y}_{cl-op}$ and Y_{op} . Then by using the associativity (1.52) and that of open-string vertex operator algebra, it is easy to show that (1.33) converges absolutely to (1.34) in the required domain.

The rest cases can all be reduced to above cases by using (1.39) (see the proof of Theorem 2.11 in [HKo2] for reference). More precisely, for $z_1^{(i)}, \dots, z_k^{(i)}, z_1, \dots, z_l \in \mathbb{H}; r_1^{(i)} > \dots > r_m^{(i)} \geq 0, r_1 > \dots > r_n > 0^4$, there always exists $a \in \mathbb{R}_+$ small enough so that both of the following sets

$$\begin{aligned} &\{z_1^{(i)} + a, \dots, z_k^{(i)} + a, r_1^{(i)} + a, \dots, r_k^{(i)} + a\}, \\ &\{z_1 - a, \dots, z_l - a, r_1 - a, \dots, r_n - a\} \end{aligned}$$

are sets whose elements have distinct absolute values. Then (1.33) equals to the following iterate series:

$$\begin{aligned} &\sum_{n_2} \sum_{n_1} m_{cl-op}^{(l;n)}(u_1, \dots, u_l; v_1, \dots, v_{i-1}, P_{n_2} e^{-aD_{op}} P_{n_1} m_{cl-op}^{(m;k)}(\tilde{u}_1, \dots, \tilde{u}_k; \\ &\quad \tilde{v}_1, \dots, \tilde{v}_m; z_1^{(i)} + a, \overline{z_1^{(i)}} + a, \dots, z_k^{(i)} + a, \overline{z_k^{(i)}} + a; r_1^{(i)} + a, \dots, r_m^{(i)} + a), \\ &\quad v_{i+1}, \dots, v_n; z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n) a_1^{n_1} a_2^{n_2} \end{aligned} \quad (1.88)$$

when $a_1 = a_2 = 1$. Hence we first switch the order of above iterate sum. Then using (1.41) and the analytic extension properties of (1.83), we can easily show that the iterate series (1.88) with opposite summing order is absolutely convergent when $1 \geq |a_1|, |a_2|$. Then we can switch the order of iterate sum in (1.88) freely without changing the value of the double sum (1.88). By (1.41), we have reduced all the remaining cases to the previous cases.

The proof of the second convergence property ((1.35) and (1.36)) is entirely the same except that one use the associativity (1.56) and (1.42). We omit the details.

It is also clear that such open-closed field algebra over V is unique because products of $\mathbb{Y}_{cl-op}$ and Y_{op} determine a dense subset of the set of all cases. The rest cases are uniquely determined by continuity. $\blacksquare$

2 Operadic formulation

In this section, we first review the notion of 2-colored partial operad and algebra over it. Then we recall the notion of Swiss-cheese partial operad $\mathfrak{S}$, its relation to sphere partial operad K and its $\mathbb{C}$ -extension $\tilde{\mathfrak{S}}^c$ [HKo1]. In the end, we prove that an open-closed field algebra over V canonically gives an algebra over $\tilde{\mathfrak{S}}^c$.

⁴If $r_n = 0$, we can further introduce another small real variable b , using (1.39) to move $r_n = 0$ to some $r'_n > 0$. We omit the detail.

2.1 2-colored partial operads

We recall the notion of 2-colored (partial) operad [V][Kt] and algebras over it. The 2-colored operad is called *relative operad* in [V] and *colored operad* in [Kt].

We first recall some basic notions from [H4]. The notion of (partial) operad can be defined in any symmetric monoidal category [MSS]. In this work, we only work in the category of sets. We will use the definition of (partial) operad given in [H4]. We denote a (partial) operad as a triple $(\mathcal{P}, I_{\mathcal{P}}, \gamma_{\mathcal{P}})$, where $\mathcal{P} = \{\mathcal{P}(n)\}_{n \in \mathbb{N}}$ is a family of sets, $I_{\mathcal{P}}$ the identity element and $\gamma_{\mathcal{P}}$ the substitution map. Note that the definition of (partial) operad in [H4] is slightly different from that in [MSS] for the appearance of $\mathcal{P}(0)$ which is very important for the study of conformal field theory. If a triple $(\mathcal{P}, I_{\mathcal{P}}, \gamma_{\mathcal{P}})$ satisfies all the axioms of a (partial) operad except the associativity of $\gamma_{\mathcal{P}}$, then it is called (*partial*) *nonassociative operad*.

We consider an important example of partial nonassociative operad. Let $U = \oplus_{n \in J} U_{(n)}$ be a vector space graded by an index set J and $E_U = \{E_U(n)\}_{n \in \mathbb{N}}$ a family of vector spaces, where

$$E_U(n) = \text{Hom}_{\mathbb{C}}(U^{\otimes n}, \overline{U}).$$

For $k, n_1, \dots, n_k \in \mathbb{N}$, $f \in E_U(k)$, $g_i \in E_U(n_i)$, $i = 1, \dots, k$ and $v_j \in U$, $j = 1, \dots, n_1 + \dots + n_k$,

$$\begin{aligned} & \gamma_{E_U}(f; g_1, \dots, g_k)(v_1 \otimes \dots \otimes v_{n_1 + \dots + n_k}) \\ &:= \sum_{s_1, \dots, s_k \in J} f(P_{s_1} g_1(v_1 \otimes \dots \otimes v_{n_1}) \otimes \dots \\ & \quad \otimes P_{s_k} g_k(v_{n_1 + \dots + n_{k-1} + 1} \otimes \dots \otimes v_{n_1 + \dots + n_k})) \end{aligned} \quad (2.1)$$

is well-defined if the sum is countable and absolutely convergent. γ_{E_U} is not associative in general because an iterate series may converge in both order but may not converge to the same value. It is clear that $(E_U, \text{id}_U, \gamma_{E_U})$ is a partial nonassociative operad. We sometimes denote it simply as E_U . If J is the set of equivalence classes of irreducible modules over a group G and $U_{(n)}$ is a direct sum of irreducible G -modules of the equivalence class $n \in J$, we denote this partial nonassociative operad as E_U^G .

Definition 2.1. An *algebra over a partial operad* $(\mathcal{P}, I_{\mathcal{P}}, \gamma_{\mathcal{P}})$, or simply a $\mathcal{P}$ -algebra, is a graded vector space U , together with a partial nonassociative operad homomorphism $\nu : \mathcal{P} \rightarrow E_U$.

Definition 2.2. Given a partial operad $(\mathcal{P}, I_{\mathcal{P}}, \gamma_{\mathcal{P}})$, a subset G of $\mathcal{P}(1)$ is called *rescaling group* for $\mathcal{P}$ if

1. For any $n \in \mathbb{N}$, $P_i \in G$, $i = 0, \dots, n$ and $P \in \mathcal{P}(n)$, $\gamma_{\mathcal{P}}(P; P_1, \dots, P_n)$ and $\gamma_{\mathcal{P}}(P_0; P)$ are well-defined.
2. $I_{\mathcal{P}} \in G$ and G together with the identity $I_{\mathcal{P}}$ and multiplication map $G \times G \xrightarrow{\gamma_{\mathcal{P}}} G$ is a group.

Definition 2.3. A partial operad $(\mathcal{P}, I_{\mathcal{P}}, \gamma_{\mathcal{P}})$ is called *G-rescalable*, if for $P_i \in \mathcal{P}(n_i)$, $i = 1, \dots, k$ and $P_0 \in \mathcal{P}(k)$, then there exists $g_i \in G$, $i = 1, \dots, k$ such that

$$\gamma_{\mathcal{P}}(\gamma_{\mathcal{P}}(P_0; g_1, \dots, g_k); P_1, \dots, P_k)$$

is well-defined.

Definition 2.4. An algebra over a G -rescalable partial operad $(\mathcal{P}, I_{\mathcal{P}}, \gamma_{\mathcal{P}})$, or a G -rescalable $\mathcal{P}$ -algebra, consists of a completely reducible G -modules $U = \bigoplus_{n \in J} U_{(n)}$, where J is the set of equivalence classes of irreducible G -modules and $U_{(n)}$ is a direct sum of irreducible G -modules of the equivalence class n , and a partial nonassociative operad homomorphism $\nu : \mathcal{P} \rightarrow E_U^G$ such that $\nu : G \rightarrow \text{End } U_{(n)}$ coincides with G -module structure on $U_{(n)}$.

We denote such algebra as (U, ν) .

For $m \in \mathbb{Z}_+$, let $m = m_1 + \dots + m_n$ be an ordered partition and $\sigma \in S_n$. The block permutation $\sigma_{(m_1, \dots, m_n)} \in S_m$ is the permutation acting on $\{1, \dots, m\}$ by permuting n intervals of lengths $m_1, \dots, m_n$ in the same way that σ permute $1, \dots, n$. Let $\sigma_i \in S_{m_i}, i = 1, \dots, n$, we view the element $(\sigma_1, \dots, \sigma_n) \in S_{m_1} \times \dots \times S_{m_n}$ naturally as an element in S_m by the canonical embedding $S_{m_1} \times \dots \times S_{m_n} \hookrightarrow S_m$.

Definition 2.5. Let $(\mathcal{Q}, I_{\mathcal{Q}}, \gamma_{\mathcal{Q}})$ be an operad. A right module over $\mathcal{Q}$, or a right $\mathcal{Q}$ -module, is a family of sets $\mathcal{P} = \{\mathcal{P}(n)\}_{n \in \mathbb{N}}$ with actions of permutation groups, equipped with maps:

$$\mathcal{P}(k) \times \mathcal{Q}(n_1) \times \dots \times \mathcal{Q}(n_l) \xrightarrow{\gamma} \mathcal{P}(n_1 + \dots + n_k)$$

such that

1. For $c \in \mathcal{P}(k)$, we have

$$\gamma(c; I_{\mathcal{Q}}, \dots, I_{\mathcal{Q}}) = c. \quad (2.2)$$

2. γ is associative. Namely, for $c \in \mathcal{P}(k)$, $d_i \in \mathcal{Q}(p_i), i = 1, \dots, k$, $e_j \in \mathcal{Q}(q_j), j = 1, \dots, p_1 + \dots + p_k$, we have

$$\gamma(\gamma(c; d_1, \dots, d_k); e_1, \dots, e_{p_1 + \dots + p_k}) = \gamma(c; f_1, \dots, f_k), \quad (2.3)$$

where

$$f_s = \gamma_{\mathcal{Q}}(d_s; e_{p_1 + \dots + p_{s-1} + 1}, \dots, e_{p_1 + \dots + p_s}).$$

3. For $c \in \mathcal{P}(k; l)$, $d_i \in \mathcal{Q}(p_i), i = 1, \dots, l$, $\sigma \in S_l$ and $\tau_j \in S_{p_j}, j = 1, \dots, l$,

$$\gamma(\sigma(c); d_1, \dots, d_l) = \sigma_{(p_1, \dots, p_l)}(\gamma(c; d_1, \dots, d_l)), \quad (2.4)$$

$$\gamma(c; \tau_1(d_1), \dots, \tau_l(d_l)) = (\tau_1, \dots, \tau_l)(\gamma(c; d_1, \dots, d_l)). \quad (2.5)$$

Homomorphisms and isomorphisms between right $\mathcal{Q}$ -modules are naturally defined.

The left module over a partial operad can be similarly defined.

Definition 2.6. A right module $\mathcal{P}$ over a partial operad $\mathcal{Q}$, or a right $\mathcal{Q}$ -module, is called G -rescalable if $\mathcal{Q}$ is G -rescalable and for any $c \in \mathcal{P}(k)$, $d_i \in \mathcal{Q}(n_i), i = 1, \dots, k$, there exist $g_i \in G, i = 1, \dots, k$ such that

$$\gamma(\gamma(c; g_1, \dots, g_k); d_1, \dots, d_k)$$

is well-defined.

Definition 2.7. A 2-colored operad consists of an operad $(\mathcal{Q}, I_{\mathcal{Q}}, \gamma_{\mathcal{Q}})$, a family of sets $\mathcal{P}(m; n)$ equipped with an $S_m \times S_n$ -action for $m, n \in \mathbb{N}$, a distinguished element $I_{\mathcal{P}} \in \mathcal{P}(1, 1)$ and substitution maps γ_1, γ_2 given as follows:

$$\begin{aligned} & \mathcal{P}(k; l) \times \mathcal{P}(m_1; n_1) \times \cdots \times \mathcal{P}(m_k; n_k) \\ & \xrightarrow{\gamma_1} \mathcal{P}(m_1 + \cdots + m_k; l + n_1 + \cdots + n_k), \\ & \mathcal{P}(k; l) \times \mathcal{Q}(p_1) \times \cdots \times \mathcal{Q}(p_l) \xrightarrow{\gamma_2} \mathcal{P}(k; p_1 + \cdots + p_l), \end{aligned} \quad (2.6)$$

satisfying the following axioms:

1. The family of sets $\mathcal{P} := \{\cup_{n \in \mathbb{N}} \mathcal{P}(m; n)\}_{m \in \mathbb{N}}$ equipped with the natural S_m -action on $\mathcal{P}(m) = \cup_{n \in \mathbb{N}} \mathcal{P}(m; n)$, together with identity element $I_{\mathcal{P}}$ and substitution maps γ_1 is an operad.
2. γ_2 gives each $\mathcal{P}(k)$ a right $\mathcal{Q}$ -module structure for $k \in \mathbb{N}$.

We denote it as $(\mathcal{P}|\mathcal{Q}, (\gamma_1, \gamma_2))$ or $\mathcal{P}$ for simplicity.

Remark 2.8. The substitution map γ_1, γ_2 can be combined into a single substitution map $\gamma = (\gamma_1, \gamma_2)$:

$$\begin{aligned} & \mathcal{P}(k; l) \times \mathcal{P}(m_1; n_1) \times \cdots \times \mathcal{P}(m_k; n_k) \times \mathcal{Q}(p_1) \times \cdots \times \mathcal{Q}(p_l) \\ & \xrightarrow{\gamma} \mathcal{P}(m_1 + \cdots + m_k; n_1 + \cdots + n_k + p_1 + \cdots + p_l). \end{aligned} \quad (2.7)$$

For this reason, we also denote $(\mathcal{P}|\mathcal{Q}, (\gamma_1, \gamma_2))$ as $(\mathcal{P}|\mathcal{Q}, \gamma)$. For some examples we encounter later, γ_1, γ_2 are often defined all together in terms of γ .

Definition 2.9. 2-colored partial operad is defined similarly as that of 2-colored operad except that γ_1, γ_2 are only partially defined and $\mathcal{Q}, \mathcal{P}$ are partial operads and (2.3) holds whenever both sides exist. If only the associativities of $\gamma_{\mathcal{Q}}, \gamma_1, \gamma_2$ do not hold, then it is called 2-colored nonassociative (partial) operad.

We give an important example of 2-colored partial nonassociative operad. Let J_1, J_2 be two index sets. $U_1 = \oplus_{n \in J_1} (U_1)_{(n)}, U_2 = \oplus_{n \in J_2} (U_2)_{(n)}$ be two graded vector spaces. Consider two families of vector spaces,

$$\begin{aligned} E_{U_2}(n) &= \text{Hom}_{\mathbb{C}}(U_2^{\otimes n}, \overline{U}_2), \\ E_{U_1|U_2}(m; n) &= \text{Hom}_{\mathbb{C}}(U_1^{\otimes m} \otimes U_2^{\otimes n}, \overline{U}_1). \end{aligned} \quad (2.8)$$

We denote both of the projection operators $U_1 \rightarrow (U_1)_{(n)}, U_2 \rightarrow (U_2)_{(n)}$ as P_n for $n \in J_1$ or J_2 . For any $f \in E_{U_1|U_2}(k; l)$, $g_i \in E_{U_1|U_2}(m_i; n_i), i = 1, \dots, k$, and $h_j \in E_{U_2}(p_j)$, $j = 1, \dots, l$, we say that

$$\begin{aligned} \Gamma(f; g_1, \dots, g_n; h_1, \dots, h_l) := & \sum_{s_1, \dots, s_k \in J_1; t_1, \dots, t_l \in J_2} f(P_{s_1} g_1(u_1^{(1)}, \dots, u_{m_1}^{(1)}, v_1^{(1)}, \dots, v_{n_1}^{(1)}), \\ & \dots, P_{s_k} g_k(u_1^{(k)}, \dots, u_{m_k}^{(k)}, v_1^{(k)}, \dots, v_{n_k}^{(k)}); \\ & P_{t_1} h_1(w_1^{(1)}, \dots, w_{n_1}^{(1)}), \dots, P_{t_l} h_l(w_1^{(l)}, \dots, w_{n_l}^{(l)})) \end{aligned}$$

where $u_j^{(i)} \in U_1, v_j^{(i)} \in U_2$, is well-defined if the multiple sum is absolutely convergent. This gives rise to a partially defined substitution map Γ :

$$E_{U_1|U_2}(k; l) \times E_{U_1|U_2}(m_1; n_1) \times \cdots \times E_{U_1|U_2}(m_k; n_k) \times E_{U_2}(p_1) \times \cdots \times E_{U_2}(p_l) \\ \xrightarrow{\Gamma} E_{U_1|U_2}(m_1 + \cdots + m_k, n_1 + \cdots + n_k + p_1 + \cdots + p_l).$$

Γ does not satisfy the associativity in general. Let $E_{U_1|U_2} = \{E_{U_1|U_2}(m; n)\}_{m, n \in \mathbb{N}}$ and $E_{U_2} = \{E_{U_2}(n)\}_{n \in \mathbb{N}}$. It is obvious that $(E_{U_1|U_2}|E_{U_2}, \Gamma)$ is a 2-colored nonassociative partial operad.

Let U_1 be a completely reducible G_1 -modules and U_2 a completely reducible G_2 -modules. Namely, $U_1 = \oplus_{n_1 \in J_1} (U_1)_{(n_1)}, U_2 = \oplus_{n_2 \in J_2} (U_2)_{(n_2)}$ where J_i is the set of equivalence classes of irreducible G_i -modules and $(U_i)_{(n_i)}$ is a direct sum of irreducible G_i -modules of the equivalence class n_i for $i = 1, 2$. In this case, we denote $E_{U_1|U_2}$ by $E_{U_1|U_2}^{G_1|G_2}$.

Definition 2.10. A homomorphism between two 2-colored (partial) operads $(\mathcal{P}_i|_{\mathcal{Q}_i}, \gamma_i), i = 1, 2$ consists of two (partial) operad homomorphisms:

$$\nu_{\mathcal{P}_1|_{\mathcal{Q}_1}} : \mathcal{P}_1 \rightarrow \mathcal{P}_2, \quad \text{and} \quad \nu_{\mathcal{Q}_1} : \mathcal{Q}_1 \rightarrow \mathcal{Q}_2$$

such that $\nu_{\mathcal{P}_1|_{\mathcal{Q}_1}} : \mathcal{P}_1 \rightarrow \mathcal{P}_2$, where $\mathcal{P}_2$ is a right $\mathcal{Q}_1$ -module by $\nu_{\mathcal{Q}_1}$, is also a right $\mathcal{Q}_1$ -module homomorphism.

Definition 2.11. An algebra over a 2-colored partial operad $(\mathcal{P}|\mathcal{Q}, \gamma)$, or a $\mathcal{P}|\mathcal{Q}$ -algebra consists of two graded vector spaces U_1, U_2 and a homomorphism $(\nu_{\mathcal{P}|\mathcal{Q}}, \nu_{\mathcal{Q}})$ from $(\mathcal{P}|\mathcal{Q}, \gamma)$ to $(E_{U_1|U_2}|E_{U_2}, \Gamma)$. We denote this algebra as $(U_1|U_2, \nu_{\mathcal{P}|\mathcal{Q}}, \nu_{\mathcal{Q}})$.

Definition 2.12. If a 2-colored partial operad $(\mathcal{P}|\mathcal{Q}, \gamma)$ is so that $\mathcal{P}$ is a G_1 -rescalable partial operad and a G_2 -rescalable right $\mathcal{Q}$ -module, then it is called $G_1|G_2$ -rescalable.

Definition 2.13. A $G_1|G_2$ -rescalable $\mathcal{P}|\mathcal{Q}$ -algebra $(U_1|U_2, \nu_{\mathcal{P}|\mathcal{Q}}, \nu_{\mathcal{Q}})$ is a $\mathcal{P}|\mathcal{Q}$ -algebra so that $\nu_{\mathcal{P}|\mathcal{Q}} : \mathcal{P} \rightarrow E_{U_1|U_2}^{G_1|G_2}$ and $\nu_{\mathcal{Q}} : \mathcal{Q} \rightarrow E_{U_2}^{G_2}$; moreover, $\nu_{\mathcal{P}|\mathcal{Q}} : G_1 \rightarrow \text{End } U_1$ coincides with the G_1 -module structure on U_1 and $\nu_{\mathcal{Q}} : G_2 \rightarrow \text{End } U_2$ coincides with the G_2 -module structure on U_2 .

2.2 Swiss-cheese partial operad $\mathfrak{S}$

A disk with strips and tubes of type $(m_-, m_+; n_-, n_+)$ ($m_-, m_+, n_-, n_+ \in \mathbb{N}$) is a disk S with the following additional data:

1. $m_- + m_+$ distinct ordered punctures $p_{-m_-}^B, \dots, p_{-1}^B, p_1^B, \dots, p_{m_+}^B$ (called *boundary punctures*) on ∂S (the boundary of S), where $p_{-m_-}^B, \dots, p_{-1}^B$ are negatively oriented and $p_1^B, \dots, p_{m_+}^B$ are positively oriented, together with local coordinates:

$$(U_{-m_-}^B, \varphi_{-m_-}^B), \dots, (U_{-1}^B, \varphi_{-1}^B); \quad (U_1^B, \varphi_1^B), \dots, (U_{m_+}^B, \varphi_{m_+}^B),$$

where U_i^B is a neighborhood of p_i^B and $\varphi_i^B : U_i^B \rightarrow \tilde{\mathbb{H}}$ is an analytic map which vanishes at p_i^B and maps $U_i^B \cap \partial S$ to $\mathbb{R}$, for each $i = -m_-, \dots, -1, 1, \dots, m_+$.

2. $n_- + n_+$ distinct ordered points $p_{-n_-}^I, \dots, p_{-1}^I, p_1^I, \dots, p_{n_+}^I$ (called *interior punctures*) in the interior of S , where $p_{-n_-}^I, \dots, p_{-1}^I$ are negatively oriented and $p_1^I, \dots, p_{n_+}^I$ are positively oriented, together with local coordinates:

$$(U_{-n_-}^I, \varphi_{-n_-}^I), \dots, (U_{-1}^I, \varphi_{-1}^I); \quad (U_1^I, \varphi_1^I), \dots, (U_{n_+}^I, \varphi_{n_+}^I),$$

where U_j^I is a local neighborhood of p_j^I and $\varphi_j^I : U_j^I \rightarrow \mathbb{C}$ is an analytic map which vanishes at p_j^I for each $j = -n_-, \dots, -1, 1, \dots, n_+$.

Two disks with strips and tubes are conformal equivalent if there exists between them a biholomorphic map which maps punctures to punctures and preserves the order of punctures and the germs of local coordinate maps. We denote the moduli space of the conformal equivalence classes of disks with strips and tubes of type $(m_-, m_+; n_-, n_+)$ as $\mathbb{S}(m_-, m_+ | n_-, n_+)$. The structure on $\mathbb{S}(m_-, m_+ | n_-, n_+)$ will be discussed in [K3]. In this work, we are only interested in disks with strips and tubes of types $(1, m_+; 0, n_+)$ for $n, l \in \mathbb{N}$. For simplicity, we denote $\mathbb{S}(1, m_+ | 0, n_+)$ by $\Upsilon(m_+; n_+)$. For such disks, we label the only negatively oriented boundary puncture as the 0-th boundary puncture.

We can choose a canonical representative for each conformal equivalence class in $\Upsilon(m_+; n_+)$ just as we did for disks with strips [HKo1] and sphere with tubes [H4]. More precisely, for a disk with strips and tubes of type $(1, m_+; 0, n_+)$ where $m_+ > 0$, we first use a conformal map f to map the disk to $\hat{\mathbb{H}}$. Then we use an automorphism of $\hat{\mathbb{H}}$ to move the only negatively oriented puncture (the 0-th puncture) to ∞ and the smallest r_k to 0, and fix the local coordinate map f_0 at ∞ to be so that $\lim_{w \rightarrow \infty} w f_0(w) = -1$. As a consequence, the canonical representative of a generic conformal equivalent class of disk with strips and tubes $Q \in \Upsilon(m_+; n_+)$ is a disk $\hat{\mathbb{H}}$, together with a negatively oriented boundary puncture at $\infty \in \hat{\mathbb{H}}$ and local coordinate map given by

$$f_0^B(w) = -e^{\sum_{j=1}^{\infty} -B_j^{(0)} w^{-j+1} \frac{d}{dw}} \frac{1}{w},$$

where $B_j^{(0)} \in \mathbb{R}$, and positively oriented boundary punctures at $r_1, \dots, r_{m_+} \in \mathbb{R}_+ \cup \{0\}$ ($r_k = 0$) and local coordinate maps given by

$$f_i^B(w) = e^{\sum_{j=1}^{\infty} B_j^{(i)} x^{j+1} \frac{d}{dx}} (b_0^{(i)})^x \frac{d}{dx} x \Big|_{x=w-r_i}, \quad i = 1, \dots, m_+,$$

where $B_j^{(i)} \in \mathbb{R}, b_0^{(i)} \in \mathbb{R}_+$, and positively oriented interior punctures at $z_1, \dots, z_{n_+} \in \mathbb{H}$ with local coordinate maps given by

$$f_i^I(w) = e^{\sum_{j=1}^{\infty} A_j^{(i)} x^{j+1} \frac{d}{dx}} (a_0^{(i)})^x \frac{d}{dx} x \Big|_{x=w-z_i}, \quad i = 1, \dots, n_+.$$

where $A_j^{(i)} \in \mathbb{C}$ and $a_0^{(i)} \in \mathbb{C}^\times$.

Let $\Pi_{\mathbb{R}} (\Pi_{\mathbb{C}})$ be the set of sequences of real numbers (complex numbers) $\{C_j\}_{j=1}^{\infty}$ such that $e^{\sum_{j>0} C_j x^{j+1} \frac{d}{dx}} x$ as a power series converges in some neighborhood of 0. We define

$$\tilde{\Lambda}^n := \{(r_1, \dots, r_n) \mid \exists \sigma \in S_n, r_{\sigma(1)} > \dots > r_{\sigma(n)} = 0\}.$$

Using the data on the canonical representative of Q , we denote $Q \in \Upsilon(m_+; n_+)$ as follows

$$[r_1, \dots, r_{m_+-1}; B^{(0)}, (b_0^{(1)}, B^{(1)}), \dots, (b_0^{(m_+)}, B^{(m_+)})] \\ [z_1, \dots, z_{n_+}; (a_0^{(1)}, A^{(1)}), \dots, (a_0^{(n_+)}, A^{(n_+)})], \quad (2.9)$$

where $(r_1, \dots, r_{m_+}) \in \tilde{\Lambda}^{m_+}$ and $b_0^{(i)} \in \mathbb{R}_+$, $a_0^{(j)} \in \mathbb{C}^\times$, and $B^{(i)} = \{B_j^{(i)}\}_{j=1}^\infty \in \Pi_{\mathbb{R}}$ and $A^{(p)} = \{A_q^{(p)}\}_{q=1}^\infty \in \Pi_{\mathbb{C}}$ for all $i = 1, \dots, m_+$, $p = 1, \dots, n_+$.

Using notations (0.1) and (0.2), for $m_+ > 0$ and $n_+ \in \mathbb{N}$, we can express the moduli space of disks with strips and tubes of type $(1, m_+; 0, n_+)$ as follows:

$$\Upsilon(m_+; n_+) = \tilde{\Lambda}^{m_+-1} \times \Pi_{\mathbb{R}} \times (\mathbb{R}_+ \times \Pi_{\mathbb{R}})^{m_+} \times M_{\mathbb{H}}^{n_+} \times (\mathbb{C}^\times \times \Pi_{\mathbb{C}})^{n_+}.$$

For $m_+ = 0, n_+ \in \mathbb{N}$, we used automorphism of $\hat{\mathbb{H}}$ to fix $B_1^{(0)} = 0$. Hence, we have

$$\Upsilon(0; n_+) = \{B^{(0)} \in \Pi \mid B_1^{(0)} = 0\} \times (\mathbb{C}^\times \times \Pi_{\mathbb{C}})^{n_+}.$$

Note that $\Upsilon(m_+; 0)$ is nothing but $\Upsilon(m_+)$ introduced and studied in [HKo1]. $\Upsilon := \{\Upsilon(n)\}_{n \in \mathbb{N}}$ is a partial operad of disks with strips [HKo1]. The identity I_Υ is an element of $\Upsilon(1; 0)$. Also, for $m_+, n_+ \in \mathbb{N}$, S_{n_+} acts on $\Upsilon(m_+; n_+)$ in the obvious way. Let $\mathfrak{S}(m) = \cup_{n_+ \in \mathbb{N}} \Upsilon(m_+; n_+)$ for $m \in \mathbb{N}$, and $\mathfrak{S} = \cup_{m \in \mathbb{N}} \mathfrak{S}(m)$.

There are two kinds of sewing operations on $\mathfrak{S}$. The first kind is called *boundary sewing operation* which sews the positively oriented boundary puncture in the first disk with a negatively oriented boundary puncture in the second disk. The second is called *interior sewing operation* which sews a positively oriented interior puncture in a disk with a negatively oriented puncture in a sphere with tubes. We describe these sewing operations more precisely. Let us consider $P \in \Upsilon(m_+; n_+)$ and $Q \in \Upsilon(p_+; q_+)$ ($Q \in K(p_+)$) for boundary sewing operations (for interior sewing operations). Let B^r ($\bar{B}^r$) denote the open (closed) ball in $\mathbb{C}$ center at 0 with radius r , φ_i the germs of local coordinate map at i -th boundary (interior) puncture p of P , and ψ_0 the germs of local coordinate map at 0-th puncture q of Q . Then we say that the i -th strip of P can be sewn with the 0-th strip (tube) of Q if there is a $r \in \mathbb{R}_+$ such that p and q are the only punctures in $\varphi_i^{-1}(\bar{B}^r)$ and $\psi_0^{-1}(\bar{B}^{1/r})$ respectively. A new disk with strips and tubes in $\Upsilon(m_+ + p_+ - 1, n_+ + q_+)$ ($\Upsilon(m_+, n_+ + p_+ - 1)$), denoted as $P_i \propto_0^B Q$ ($P_i \propto_0^I Q$), is obtained by cutting out $\varphi_i^{-1}(B^r)$ and $\psi_0^{-1}(B^{1/r})$ from P and Q respectively, and then identifying the boundary of $\varphi_i^{-1}(\bar{B}^r)$ and $\psi_0^{-1}(\bar{B}^{1/r})$ via the map

$$\psi^{-1} \circ J_{\hat{\mathbb{H}}} \circ \varphi_i$$

where $J_{\hat{\mathbb{H}}} : w \rightarrow -\frac{1}{w}$.

The boundary sewing operations and interior sewing operations induce the following partially defined substitution maps:

$$\begin{aligned} \Upsilon(k; l) \times \Upsilon(m_1; n_1) \times \dots \times \Upsilon(m_k; n_k) \times K(p_1) \times \dots \times K(p_l) \\ \xrightarrow{\gamma} \Upsilon(m_1 + \dots + m_k; n_1 + \dots + n_k + p_1 + \dots + p_l). \end{aligned} \quad (2.10)$$

The following Proposition is clear.

Proposition 2.14. $(\mathfrak{S}|K, \gamma)$ is a $\mathbb{R}_+|\mathbb{C}^\times$ -rescalable 2-colored partial operad.

This $\mathbb{R}_+|\mathbb{C}^\times$ -rescalable 2-colored partial operad $(\mathfrak{S}|K, \gamma)$ is a generalization of Voronov's Swiss-cheese operad [V]. So we will call it *Swiss-cheese partial operad* and sometimes denote it by $\mathfrak{S}$ for simplicity. The relation between Swiss-cheese operad and Swiss-cheese partial operad is an analogue of that between little disk operad and sphere partial operad [H4].

2.3 Sewing equations and the doubling map δ

We are interested in finding the canonical representative of disk with strips and tubes obtained by sewing two such disks or sewing a disk with a sphere. In the case of sphere partial operad K , such canonical representatives were obtained by Huang [H1][H2][H4] by solving the so-called sewing equation. Similarly, the canonical representatives of disks with strips and tubes obtained by two types of sewing operations can also be determined by solving two types of sewing equations.

We start with the boundary sewing operations. For $Q \in \mathfrak{S}$, we denoted the canonical representative of Q as Σ_Q . Let $P \in \Upsilon(m; n)$ and $Q \in \Upsilon(p; q)$. Let g_0 be the local coordinate map at $\infty \in \Sigma_Q$ and f_i be that at $z_i \in \Sigma_P, 1 \leq i \leq m$. We assume that $P_i \infty_0^B Q$ exists. Then the canonical disk $\Sigma_{P_i \infty_0^B Q}$ can be obtained by solving the following sewing equation:

$$F_{(1)}^B(w) = F_{(2)}^B \left(g_0^{-1} \left(\frac{-1}{f_i(w)} \right) \right) \quad (2.11)$$

where $F_{(1)}^B$ is a conformal map from an open neighborhood of $\infty \in \Sigma_P$ to an open neighborhood of $\infty \in \Sigma_{P_i \infty_0^B Q}$, and $F_{(2)}^B$ is a conformal map from an open neighborhood of $0 \in \Sigma_Q$ to an open neighborhood of $0 \in \Sigma_{P_i \infty_0^B Q}$, with the following normalization conditions:

$$F_{(1)}^B(\infty) = \infty, \quad F_{(2)}^B(0) = 0, \quad \lim_{w \rightarrow \infty} \frac{F_{(1)}^B(w)}{w} = 1. \quad (2.12)$$

It is easy to see that the solution of (2.11) and (2.12) is unique. Notice that $F_{(1)}^B, F_{(2)}^B, f_0$ and f_i in (2.11) are all real analytic.

Similarly, let $P \in \Upsilon(m; n)$ and $Q \in K(p)$. We denote the canonical sphere with tubes of Q as Σ_Q . Let g_0 be the local coordinate map at $\infty \in Q$ and f_i be that at $z_i \in P, 1 \leq i \leq n$. We assume that $P_i \infty_0^I Q$ exists. Then $\Sigma_{P_i \infty_0^I Q}$ can be obtained by solving the following sewing equation:

$$F_{(1)}^I(w) = F_{(2)}^I \left(g_0^{-1} \left(\frac{-1}{f_i(w)} \right) \right) \quad (2.13)$$

where $F_{(1)}^I$ is a conformal map from an open neighborhood of $\hat{\mathbb{R}}$ in Σ_P to an open neighborhood of $\hat{\mathbb{R}}$ in $\Sigma_{P_i \infty_0^I Q}$, and $F_{(2)}^I$ is a conformal map from an open neighborhood of 0 in Σ_Q to an open subset of $\mathbb{H} \subset \Sigma_{P_i \infty_0^I Q}$, with the following normalization conditions:

$$F_{(1)}^I(\infty) = \infty, \quad F_{(2)}^I(0) = 0, \quad \lim_{w \rightarrow \infty} \frac{F_{(1)}^I(w)}{w} = 1. \quad (2.14)$$

It is easy to see that the solution of (2.13) and (2.14) is unique as well.

Notice that $F_{(1)}^I$ is real analytic because it maps $\mathbb{R}$ to $\mathbb{R}$. Hence, $F_{(1)}^I$ is also the unique solution for the following equation

$$F_{(1)}^I(w) = \overline{F_{(2)}^I} \left(\overline{g_0^{-1}} \left(\frac{-1}{\overline{f_i(w)}} \right) \right) \quad (2.15)$$

with the same normalization condition (2.14).

In sphere partial operad, we define a complex conjugation map $\text{Conj} : K \rightarrow K$ as follows

$$\begin{aligned} \text{Conj} : (z_1, \dots, z_{n-1}; A^{(0)}, (a_0^{(1)}, A^{(1)}), \dots, (a_0^{(n)}, A^{(n)})) \\ \longmapsto (\bar{z}_1, \dots, \bar{z}_{n-1}; \bar{A}^{(0)}, (\bar{a}_0^{(1)}, \bar{A}^{(1)}), \dots, (\bar{a}_0^{(n)}, \bar{A}^{(n)})). \end{aligned} \quad (2.16)$$

For simplicity, we denote $\text{Conj}(Q)$ as $\bar{Q}$ for $Q \in K$.

Proposition 2.15. *Conj is an partial operad automorphism of K .*

Proof. It is clear that $\text{Conj}((\mathbf{0}, (1, \mathbf{0}))) = (\mathbf{0}, (1, \mathbf{0}))$ and Conj is equivariant with respect to the action of permutation group. Moreover, Conj is obviously bijective. It only remains to show that, for $1 \leq i \leq m$,

$$\overline{P_i \infty_0 Q} = \bar{P}_i \infty_0 \bar{Q} \quad (2.17)$$

for any pair $P \in K(m), Q \in K(n)$ such that $P_i \infty_0 Q$ exists.

Let f_i be the local coordinate map at i -th puncture in P and g_0 be that at ∞ in Q . Then the local coordinate map at i -th puncture in $\bar{P}$ is $\bar{f}_i$ and that at ∞ in $\bar{Q}$ is $\bar{g}_0$. Also notice that the sewing equation and normalization equation for sphere partial operad ([H4]) are the same as the equation (2.11) and (2.12). Let $F_{\text{sphere}}^{(1)}, F_{\text{sphere}}^{(2)}$ be the solution of (2.11) and (2.12) for the sphere with tubes, then $\bar{F}_{\text{sphere}}^{(1)}, \bar{F}_{\text{sphere}}^{(2)}$ also satisfy the same normalization condition and the following sewing equation

$$\bar{F}_{\text{sphere}}^{(1)}(w) = \bar{F}_{\text{sphere}}^{(2)}\left(\overline{g_0^{-1}}\left(\frac{-1}{\bar{f}_i(w)}\right)\right),$$

which is the sewing equation for $\bar{P}_i \infty_0 \bar{Q}$. Using the explicit formula ((A.6.1)-(A.6.5) in [H4]) of the moduli $\bar{P}_i \infty_0 \bar{Q}$ in terms of $\bar{F}_{\text{sphere}}^{(1)}, \bar{F}_{\text{sphere}}^{(2)}, \bar{f}_i, \bar{g}_0$, one can easily see that (2.17) is true. $\blacksquare$

There is a canonical doubling map $\delta : \mathfrak{S} \rightarrow K$ defined as follows. Let $Q \in \Upsilon(n; l)$ with form

$$Q = [r_1, \dots, r_{n-1}; B^{(0)}, (b_0^{(1)}, B^{(1)}), \dots, (b_0^{(n)}, B^{(n)}) | z_1, \dots, z_l; (a_0^{(1)}, A^{(1)}), \dots, (a_0^{(l)}, A^{(l)})].$$

Then

$$\begin{aligned} \delta(Q) = (z_1, \dots, z_l, \bar{z}_1, \dots, \bar{z}_l, r_1, \dots, r_{n-1}; (a_0^{(1)}, A^{(1)}), \dots, (a_0^{(l)}, A^{(l)}), \\ (\overline{a_0^{(1)}}), \overline{A^{(1)}}), \dots, (\overline{a_0^{(l)}}), \overline{A^{(l)}}); B^{(0)}, (b_0^{(1)}, B^{(1)}), \dots, (b_0^{(n)}, B^{(n)})). \end{aligned} \quad (2.18)$$

Proposition 2.16. *Let $P \in \Upsilon(m; n)$ and $Q \in \Upsilon(p; q)$. Assume that $P_i \infty_0^B Q$ exists. We have*

$$\delta(P_i \infty_0^B Q) = \delta(P)_{2n+i} \infty_0 \delta(Q). \quad (2.19)$$

Proof. Since $F_{(1)}^B, F_{(2)}^B, g_0^{-1}$ and f_i in (2.11) are all real analytic, every solution of (2.11) and (2.12) for disk with strips and tubes can be extended to a solution of the same sewing equation for sphere with tubes by Schwarz's reflection principle. Then the Proposition follows immediately from this fact. $\blacksquare$

Given a canonical disk with strips and tubes Σ_Q corresponding to moduli $Q \in \mathfrak{S}$, we consider its complex conjugation, denoted as $\overline{\Sigma_Q}$. $\overline{\Sigma_Q}$ is the lower half plane $\hat{\mathbb{H}}$ together with the same boundary punctures and local coordinate maps as those in Σ_Q , and interior punctures which are the complex conjugation of the interior punctures in Σ_Q with local coordinate maps being the complex conjugation of those in Σ_Q . We can denote it as

$$[r_1, \dots, r_{m_+-1}; B^{(0)}, (b_0^{(1)}, B^{(1)}), \dots, (b_0^{(m_+)}, B^{(m_+)}) | \bar{z}_1, \dots, \bar{z}_{n_+}; (\overline{a_0^{(1)}}, \overline{A^{(1)}}), \dots, (\overline{a_0^{(n_+)}}), \overline{A^{(n_+)}})] \quad (2.20)$$

where $z_1, \dots, z_{n_+} \in \mathbb{H}$.

Lemma 2.17. *Let $P \in \Upsilon(m; n)$ and $Q \in K(p)$. Assume that $P_i \infty_0^I Q$ exists for $1 \leq i \leq m$. Then $\overline{\Sigma_{P_i} \infty_0^I \bar{Q}}$ also exists and*

$$\overline{\Sigma_{P_i} \infty_0^I Q} \cong \overline{\Sigma_P} \infty_0^I \bar{Q}. \quad (2.21)$$

Proof. We can choose a canonical representative of $\overline{\Sigma_P} \infty_0^I \bar{Q}$ as a lower half plane $\hat{\mathbb{H}}$ with two punctures at ∞ and 0 and the local coordinate map g_0 at ∞ being so that $\lim_{w \rightarrow \infty} w g_0(w) = -1$. We denote such representative of $\overline{\Sigma_P} \infty_0^I \bar{Q}$ (a lower half plane) as Σ_1 . Σ_1 can be obtained by solving the sewing equation

$$G_{(1)}^I(w) = G_{(2)}^I \left(\overline{g_0^{-1}} \left(\frac{-1}{\overline{f_i(w)}} \right) \right), \quad (2.22)$$

where $G_{(1)}^I$ is a conformal map from a neighborhood of $\hat{\mathbb{R}} \subset \overline{\Sigma_P}$ to a neighborhood of $\hat{\mathbb{R}} \subset \Sigma_1$, and $G_{(2)}^I$ is a conformal map from a neighborhood of $0 \in \overline{\Sigma_Q}$ to an open subset of $\mathbb{H} \subset \Sigma_1$, with the normalization equation (2.14). By comparing (2.22) with (2.15), we see that the unique solution $F_{(1)}^I$ (real analytic) and $\overline{F_{(2)}^I}$ of (2.15) and (2.14) exactly gives the unique solution $G_{(1)}^I$ and $G_{(2)}^I$ of (2.22) and (2.14). Hence we have $\Sigma_1 = \overline{\Sigma_{P_i} \infty_0^I Q}$. ■

Proposition 2.18. *Let $P \in \Upsilon(n; l)$, $Q \in K(m)$ and $P_i \infty_0^I Q$ exists for $1 \leq i \leq l$. Then*

$$\delta(P_i \infty_0^I Q) = (\delta(P)_{i \infty_0} Q)_{l+m-1+i \infty_0} \bar{Q} \quad (2.23)$$

Proof. Now we first consider the right hand side of (2.23). We denote the canonical representative of any $R \in K(l)$ as Σ_R . Then $\Sigma_{\delta(P)}$ can be viewed as a union of the closure of upper half plane, denoted as U_+ , and the closure of lower half plane, denoted as U_- . Let Σ_+ be the Riemann surface obtained by sewing U_+ with Q , and Σ_- the Riemann surface obtained by sewing U_- with $\bar{Q}$. By identifying the real line in $U_+ \subset \Sigma_+$ with the real line in $U_- \subset \Sigma_-$ using identity map, we obtain the surface $\Sigma_+ \# \Sigma_-$ which is isomorphic to the canonical sphere with tubes $\Sigma_{(\delta(P)_i \infty_0 Q)_{l+m-1+i \infty_0} \bar{Q}}$.

Both Σ_+ and Σ_- are disks with strips and tubes. Since $U_+ = \Sigma_P$, there is a unique biholomorphic map f from Σ_+ to the canonical disk with strips and tubes $\Sigma_{Q_i \infty_0^I P}$. Similarly, because $U_- = \overline{\Sigma_P}$, by Lemma 2.17, there is a unique biholomorphic map g from Σ_- to the canonical disk with strips and tubes $\overline{\Sigma_{P_i \infty_0^I Q}}$. The restriction of f on the neighborhood of $\hat{\mathbb{R}} = \partial U_+$ is nothing but the unique real analytic map $F_{(1)}^I$ satisfying

(2.13) and (2.14). Meanwhile, the restriction of g on a neighborhood of $\hat{\mathbb{R}} = \partial U_-$ is nothing but the same $F_{(1)}^I$ satisfying (2.15) and (2.14). So we must have $f|_{\hat{\mathbb{R}}} = g|_{\hat{\mathbb{R}}}$, which further implies that $f^{-1}|_{\hat{\mathbb{R}}} = g^{-1}|_{\hat{\mathbb{R}}}$. Hence, f^{-1} can be extended to a biholomorphic map from $\Sigma_{\delta(P_i \infty_0^I Q)}$ to $\Sigma_+ \# \Sigma_-$.

Therefore $\Sigma_{\delta(P_i \infty_0^I Q)}$ must be biholomorphic to $\Sigma_{(\delta(P)_i \infty_0 Q)_{l+m-1+i} \infty_0 \bar{Q}}$. Since they are both canonical representatives of sphere with tubes, we must have the equality (2.23). $\blacksquare$

Remark 2.19. Proposition 2.18 is nothing but the doubling trick [A][C1] stated in our partial operad language. It also implies that the bulk theories in an open-closed conformal field theory must contain both chiral parts and anti-chiral parts.

Corollary 2.20. *For $P \in \delta(\Upsilon(n, l)) \subset K(n + 2l)$ as in (2.18) and $Q \in K(m)$, the sewing operations: $(P_i \infty_0 Q)_{l+m-1+i} \infty_0 \bar{Q}$ for $1 \leq i \leq l$ define an action of the diagonal $\{(Q, \bar{Q}) \in K \times K\}$ of $K \times K$ on $\delta(\mathfrak{S})$.*

2.4 The $\mathbb{C}$ -extensions of $\mathfrak{S}$

In order to study open-closed conformal field theories with nontrivial central charges, we need study the $\mathbb{C}$ -extensions of Swiss-cheese partial operad $\mathfrak{S}$.

For $c \in \mathbb{C}$, let $\tilde{K}^c$ be the $\frac{c}{2}$ -th power of determinant line bundle over K [H4]. We denote the pullback line bundle over $\mathfrak{S}$ through the doubling map δ as $\tilde{\mathfrak{S}}^c$. δ can certainly be extended to a map on $\tilde{\mathfrak{S}}^c$. We still denote it as δ . For any $n \in \mathbb{N}$, the restrictions of the sections ψ_{n+2l} of $\tilde{K}^c(n + 2l)$ for $l \in \mathbb{N}$ to $\Upsilon(n; l)$ gives a section of $\tilde{\mathfrak{S}}^c(n)$ and we shall use $\psi_n^{\mathfrak{S}}$ to denote this section. It is clear that $\tilde{\Upsilon}^c$, the $\mathbb{C}$ -extension of the partial operad of disks with strips, is the pullback bundle of the inclusion map $\Upsilon \hookrightarrow \mathfrak{S}$.

The boundary sewing operations in $\tilde{\mathfrak{S}}^c$ are naturally induced from the sewing operations of $\tilde{K}^c$. We denote the boundary sewing operations in $\tilde{\mathfrak{S}}^c$ as ${}_i \widetilde{\infty}_0^B$.⁵ More explicitly, let $P \in \Upsilon(n; l)$ and $Q \in \Upsilon(m; k)$ be so that $P_i \infty_0^B Q$ exists. Let $\tilde{P}, \tilde{Q}$ be elements in the fiber over P and Q respectively. We define

$$\tilde{P}_i \widetilde{\infty}_0^B \tilde{Q} := \delta^{-1}(\delta(\tilde{Q})_{2l+i} \widetilde{\infty}_0 \tilde{Q}), \quad (2.24)$$

where δ^{-1} is defined on the image of δ .

We would also like to lift an interior sewing operation ${}_i \infty_0^I$ to a sewing operation between an element in $\tilde{\mathfrak{S}}^c$ and an element in $\tilde{K}^c \otimes \overline{\tilde{K}^c}$. We still call it interior sewing operation and denote it as ${}_i \widetilde{\infty}_0^I$. Let $P \in \Upsilon(n; l), Q \in K(m)$ such that $P_i \infty_0^I Q$ exists. Let $\tilde{P}, \tilde{Q}$ be elements in the fibers over P and Q respectively. Let $\psi_m \otimes \bar{\psi}_m$ be the canonical section on $\tilde{K}^c \otimes \overline{\tilde{K}^c}(m)$. Then we have $\tilde{Q} = \lambda \psi_m \otimes \bar{\psi}_m(Q)$ for some $\lambda \in \mathbb{C}$. Then we define $\tilde{P}_i \widetilde{\infty}_0^I \tilde{Q}$ by

$$\tilde{P}_i \widetilde{\infty}_0^I \tilde{Q} := \delta^{-1}((\delta(\tilde{P})_i \widetilde{\infty}_0 \lambda \psi_m(Q))_{l+m-1+i} \widetilde{\infty}_0 \psi_m(\bar{Q})) \quad (2.25)$$

The following Lemma shows that the interior sewing operations are associative.

⁵Since we always work with a fixed $c \in \mathbb{C}$, it is convenient to make the dependence on c implicit in some notations.

Lemma 2.21. *Let $P \in \Upsilon(n; l)$, $Q_1 \in K(m_1)$, $Q_2 \in K(m_2)$ and $\tilde{P}, \tilde{Q}_1, \tilde{Q}_2$ be elements in fibers of line bundles $\tilde{\mathfrak{S}}^c$ and $\tilde{K}^c \otimes \overline{\tilde{K}^c}$ over the base points P, Q_1, Q_2 respectively. Let $1 \leq i \leq l$ and $1 \leq j \leq m_1$. Then we have*

$$(\tilde{P}_i \widetilde{\infty}_0^I \tilde{Q}_{j-1})_{i+j-1} \widetilde{\infty}_0^I \tilde{Q}_2 = \tilde{P}_i \widetilde{\infty}_0^I (\tilde{Q}_{j-1} \widetilde{\infty}_0 \tilde{Q}_2) \quad (2.26)$$

assuming that the sewing operations appeared in (2.26) are all well-defined.

Proof. Let $\lambda_i \in \mathbb{C}, i = 1, 2$ be such that $\bar{Q}_i = \lambda_i \psi_{m_i} \otimes \bar{\psi}_{m_i}(Q_i), i = 1, 2$. Let $(a_0^{(i)}, A^{(i)})$ be the local coordinate map at j -th puncture of Q_1 and Let $B^{(0)}$ be the local coordinate map at ∞ in Q_2 . By (2.25), the δ image of the left hand side of (2.26) equals to

$$((\delta(\tilde{P})_i \widetilde{\infty}_0 \lambda_1 \psi_{m_1}(Q_1))_{l+m_1-1+i} \widetilde{\infty}_0 \psi_{m_1}(\bar{Q}_1))_{i+j-1} \widetilde{\infty}_0 \lambda_2 \psi_{m_2}(Q_2)_{l+m_1+m_2+i+j-3} \widetilde{\infty}_0 \psi_{m_2}(\bar{Q}_2)$$

By the associativity of the partial operad $\tilde{K}^c$, the above formula equals to

$$\begin{aligned}
& (\delta(\tilde{P})_i \widetilde{\infty}_0 (\lambda_1 \psi_{m_1} (Q_1)_j \widetilde{\infty}_0 \lambda_2 \psi_{m_2} (Q_2))) \\
& \quad \quad \quad {}_{l+m_1+m_2-2+i} \widetilde{\infty}_0 (\psi_{m_1} (\bar{Q}_1)_j \widetilde{\infty}_0 \psi_{m_2} (\bar{Q}_2)) \\
& = (\delta(\tilde{P})_i \widetilde{\infty}_0 (\lambda_1 \lambda_2 e^{\Gamma(A^{(i)}, B^{(0)}, a_0^{(i)})c} \psi_{m_1+m_2-1} (Q_1 \text{ } j \infty_0 Q_2))) \\
& \quad \quad \quad {}_{l+m_1+m_2-2+i} \widetilde{\infty}_0 e^{\Gamma(\overline{A^{(i)}}, \overline{B^{(0)}}, \overline{a_0^{(i)}})c} \psi_{m_1+m_2-1} (\overline{Q_1 \text{ } j \infty_0 Q_2}) \\
& = \delta(\tilde{P}_i \widetilde{\infty}_0^I (\lambda_1 \lambda_2 e^{\Gamma(A^{(i)}, B^{(0)}, a_0^{(i)})c} e^{\Gamma(\overline{A^{(i)}}, \overline{B^{(0)}}, \overline{a_0^{(i)}})c} \\
& \quad \quad \quad \psi_{m_1+m_2-1} \otimes \bar{\psi}_{m_1+m_2-1} (Q_1 \text{ } j \infty_0 Q_2))) \\
& = \delta(\tilde{P}_i \widetilde{\infty}_0^I (\lambda_1 \psi_{m_1} \otimes \bar{\psi}_{m_1} (Q_1)_j \widetilde{\infty}_0 \lambda_2 \psi_{m_2} \otimes \bar{\psi}_{m_2} (Q_2))) \\
& = \delta(\tilde{P}_i \widetilde{\infty}_0^I (\tilde{Q}_1 \text{ } i \widetilde{\infty}_0 \tilde{Q}_2)),
\end{aligned}$$

which is nothing but the right hand side of (2.26).

Boundary sewing operations and interior sewing operations induce the following partially defined substitution maps $\tilde{\gamma}$:

$$\begin{aligned} & \tilde{\Upsilon}^c(n; l) \times \tilde{\Upsilon}^c(m_1; k_1) \times \cdots \times \tilde{\Upsilon}^c(m_n; k_n) \times \tilde{K}^c(n_1) \times \cdots \times \tilde{K}^c(n_l) \\ & \xrightarrow{\tilde{\gamma}} \tilde{\Upsilon}^c(m_1 + \cdots + m_n; k_1 + \cdots + k_n + n_1 + \cdots + n_l) \end{aligned}$$

The following Proposition is clear.

Proposition 2.22. $(\tilde{\mathfrak{S}}^c | \tilde{K}^c \otimes \overline{\tilde{K}^c}, \tilde{\gamma})$ is a $\mathbb{R}_+ |\mathbb{C}^\times$ -rescalable 2-colored partial operad.

We will call $(\tilde{\mathfrak{S}}^c | \tilde{K}^c \otimes \overline{\tilde{K}^c}, \tilde{\gamma})$ *Swiss-cheese partial operad with central charge c* . Note that $\tilde{\mathfrak{S}}^c$ restricted on Υ is just $\tilde{\Upsilon}^c$ which was introduced in [HKo1].

2.5 Smooth $\tilde{\mathfrak{S}}^c|_{\tilde{K}^c} \otimes \overline{\tilde{K}^c}$ -algebras

Let $V^O = \oplus_{n \in \mathbb{R}} V_{(n)}^O$, where $V_{(n)}^O$ has a structure of irreducible $\mathbb{R}_+$ -module given by $r \mapsto r^n \text{id}_{V_{(n)}^O}$ for $r \in \mathbb{R}_+$. Let $V^C = \oplus_{(m,n) \in \mathbb{R} \times \mathbb{R}} V_{(m,n)}^C$, where $V_{(m,n)}^C$ has a structure of

irreducible $\mathbb{C}^\times$ -module given by $z \mapsto z^m \bar{z}^n \text{id}_{V_{(m,n)}^C}$ for $z \in \mathbb{C}^\times$ (recall (1.16) and (1.17)) for all m, n . This also implies that $V_{(m,n)}^C = 0$ for all $m - n \notin \mathbb{Z}$. Let

$$(V^O|V^C, \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}, \nu_{\tilde{K}^c \otimes \overline{\tilde{K}^c}})$$

be a $\mathbb{R}_+|\mathbb{C}^\times$ -rescalable $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra.

Definition 2.23. The $\mathbb{R}_+|\mathbb{C}^\times$ -rescalable $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra $(V^O|V^C, \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}, \nu_{\tilde{K}^c \otimes \overline{\tilde{K}^c}})$ is called *smooth* if it satisfies the following two conditions:

1. $\dim V_{(s)}^O < \infty$ for $s \in \mathbb{R}$, $V_{(n)}^O = 0$ for $n < 0$ and $\dim V_{(m,n)}^C < \infty$ for $m, n \in \mathbb{Z}$, $V_{(m,n)}^C$ for $m < 0$ or $n < 0$.
2. For $w \in (V^O)'$, $v_1, \dots, v_m \in V^O$, $u_1, \dots, u_n \in V^C$, and $\tilde{P} \in \tilde{\Upsilon}^c(m; n)$, the following map

$$\tilde{P} \mapsto \langle w, \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}(\tilde{P})(v_1 \otimes \dots \otimes v_m \otimes u_1 \otimes \dots \otimes u_n) \rangle$$

is linear on fiber and smooth on the base space $\Upsilon(m; n)$.

By [H4], a vertex operator algebra U canonically gives a $\tilde{K}^c$ -algebra. Using the doubling map δ , this $\tilde{K}^c$ -algebra naturally gives a smooth $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra, in which $V^O = U$ and $V^C = U \otimes U$. It is also easy to see that this $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra canonically gives an analytic open-closed field algebra which is nothing but $(U \otimes U, U, \text{id}_U, \text{id}_U)$ or simply $(U \otimes U, U)$ discussed in Section 1. We still denote this smooth $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra as $(U \otimes U, U)$. A smooth $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra containing $(U \otimes U, U)$ as a subalgebra is called a *smooth $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra over U* .

Let $(V_{cl}, V_{op}, m_{cl-op})$ be an open-closed field algebra over V . By definition, V_{cl} is a conformal full field algebra over $V \otimes V$. By the results in [K2], V_{cl} has a structure of smooth $\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra structure, we denote it as $(V_{cl}, \nu_{\tilde{K}^c \otimes \overline{\tilde{K}^c}})$.

Let $Q \in \Upsilon(n; l)$ of form (2.9) such that $r_1 > \dots > r_{n-1} > r_n = 0$. We define a map $\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}} : \tilde{\mathfrak{S}}^c \rightarrow E_{V_{op}|V_{cl}}^{\mathbb{R}_+|\mathbb{C}^\times}$ as follows:

$$\begin{aligned} & \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}(\lambda \psi_n^{\mathfrak{S}}(Q))(v_1 \otimes \dots \otimes v_n \otimes u_1 \otimes \dots \otimes u_l) \\ & := \lambda e^{-L_-(B^{(0)})} m_{cl-op}^{(l;n)}(e^{-L_+(A^{(1)})}(a_0^{(1)})^{-L(0)} \otimes e^{-L_+(\overline{A^{(1)}})}(\overline{a_0^{(1)}})^{-L(0)} u_1, \\ & \quad \dots e^{-L_+(A^{(l)})}(a_0^{(l)})^{-L(0)} \otimes e^{-L_+(\overline{A^{(l)}})}(\overline{a_0^{(l)}})^{-L(0)} u_l; \\ & \quad e^{-L_+(B^{(1)})}(b_0^{(1)})^{-L(0)} v_1, \dots, e^{-L_+(B^{(n)})}(b_0^{(n)})^{-L(0)} v_n; \\ & \quad z_1, \bar{z}_1, \dots, z_l, \bar{z}_l; r_1, \dots, r_n), \end{aligned} \tag{2.27}$$

where $L_\pm(A) = \sum_{j=1}^\infty L(\pm j)A_j$ for any $A = \{A_1, A_2, \dots\}$, $A_j \in \mathbb{C}$, for $u_1, \dots, u_l \in V_{cl}$, $v_1, \dots, v_n \in V_{op}$. Let e_l be the identity element of S_l . $\forall \sigma \in S_n$, we define

$$\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}((\sigma, e_l)(\lambda \psi_n^{\mathfrak{S}}(Q))) = \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}(\lambda \psi_n^{\mathfrak{S}}(Q)). \tag{2.28}$$

We have finished the definition of $\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}$ in all cases. By results in [HKo1], the restriction of $\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}$ on $\tilde{\Upsilon}^c$ clearly gives a morphism of partial nonassociative operad from $\tilde{\Upsilon}^c$ to $E_{V_{op}}^{\mathbb{R}_+}$.

Theorem 2.24. $(V_{op}|V_{cl}, \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}, \nu_{\tilde{K}^c \otimes \overline{\tilde{K}^c}})$ is a smooth $\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}$ -algebra.

Proof. By the permutation property of open-closed field algebra and (2.28), it is clear that $\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}$ is equivariant with the actions of permutation groups.

The conditions in Definition 2.23 are automatically satisfied. It remains to show that

$$\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}} \circ \tilde{\gamma} = \Gamma \circ (\nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}, \dots, \nu_{\tilde{\mathfrak{S}}^c|\tilde{K}^c \otimes \overline{\tilde{K}^c}}, \nu_{\tilde{K}^c \otimes \overline{\tilde{K}^c}}, \dots, \nu_{\tilde{K}^c \otimes \overline{\tilde{K}^c}}) \quad (2.29)$$

as a map

$$\begin{aligned} \tilde{\Upsilon}^c(n; l) \times \tilde{\Upsilon}^c(m_1; k_1) \times \dots \times \tilde{\Upsilon}^c(m_n; k_n) \times \tilde{K}^c(n_1) \times \dots \times \tilde{K}^c(n_l) \\ \rightarrow \text{Hom}(V_{cl}^{\otimes k_1 + \dots + k_n + n_1 + \dots + n_l} \otimes V_{op}^{\otimes n + m_1 + \dots + m_n}, \overline{V_{op}}). \end{aligned} \quad (2.30)$$

Thanks to the doubling map δ and (2.26), $\tilde{\mathfrak{S}}^c$ can be viewed as a partial suboperad of $\tilde{K}^c$ with single-sewing operations for the punctures in $\mathbb{R}_+$ and double-sewing operations for mirror pairs of punctures in upper and lower half planes. By using the V -invariant boundary condition, the chirality splitting property and the convergence and extension properties of any products and iterates of intertwining operators proved by Huang for any V satisfying the condition in Theorem 0.1, it is easy to generalize the proof of Huang's fundamental result Proposition 5.4.1 in [H4] and results in [H5][H6] to show that (2.29) holds. The arguments are standard but tedious. We omit the details. ■

3 Categorical formulation

In this section, we study open-closed field algebras over V from a tensor-categorical point of view. In Section 3.1, we recall some basic ingredients of the vertex tensor categories. In Section 3.2, we reformulate the notion of open-closed field algebra over V categorically by a categorical notion called open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebra.

3.1 Vertex tensor categories

The theory of tensor products for modules over a vertex operator algebra was developed by Huang and Lepowsky [HL2]-[HL5][H3]. By Theorem 0.1 and our assumption on V , the category of V -modules, denoted as $\mathcal{C}_V$, have a structure of vertex tensor category [HL2]. In particular, it has a structure of semisimple braided tensor category.

We review some of the ingredients of vertex tensor category $\mathcal{C}_V$ and set our notations along the way.

There is a tensor product bifunctor $\boxtimes_{P(z)} : \mathcal{C}_V \times \mathcal{C}_V \rightarrow \mathcal{C}_V$ for each $P(z), z \in \mathbb{C}^\times$ in sphere partial operad K , where $P(z)$ is the conformal equivalence class of sphere with three punctures $0, z, \infty$ and standard local coordinates [H4]. We denote $\boxtimes_{P(1)}$ simply as $\boxtimes$. For any pair of V -modules W_1, W_2 , the module $W_1 \boxtimes_{P(z)} W_2$ is spanned by the homogeneous components of $w_1 \boxtimes_{P(z)} w_2 \in \overline{W_1 \otimes W_2}, \forall w_1 \in W_1, w_2 \in W_2$.

For each V -module W , there is a left unit isomorphism $l_W : V \boxtimes W \rightarrow W$ defined by

$$\overline{l_W}(v \boxtimes w) = Y_W(v, 1)w, \quad \forall v \in V, w \in W, \quad (3.1)$$

where $\overline{l_W}$ is the unique extension of l_W on $\overline{V \boxtimes W}$ and Y_W is the vertex operator which defines the module structure on W , and a right unit isomorphism $r_W : W \boxtimes V \rightarrow W$ defined by

$$\overline{r_W}(w \boxtimes v) = e^{L(-1)}Y_W(v, -1)w, \quad \forall v \in V, w \in W. \quad (3.2)$$

Remark 3.1. We have used “overline” for the extensions of maps, algebraic completions of graded vector spaces and complex conjugations of complex variables. One shall not confuse them because they acts on different things.

Let W_1 and W_2 be V -modules. For a given path $\gamma \in \mathbb{C}^\times$ from a point z_1 to z_2 , there is a parallel isomorphism associated to this path

$$\mathcal{T}_\gamma : W_1 \boxtimes_{P(z_1)} W_2 \longrightarrow W_1 \boxtimes_{P(z_2)} W_2.$$

Let $\mathcal{Y}$ be the intertwining operator corresponding to the intertwining map $\boxtimes_{P(z_2)}$ and $l(z_1)$ the value of the logarithm of z_1 determined by $\log z_2$ and analytic continuation along the path γ . For $w_1 \in W_1, w_2 \in W_2$, $\mathcal{T}_\gamma$ is defined by

$$\overline{\mathcal{T}_\gamma}(w_1 \boxtimes_{P(z_1)} w_2) = \mathcal{Y}(w_1, e^{l(z_1)})w_2,$$

where $\overline{\mathcal{T}_\gamma}$ is the natural extension of $\mathcal{T}_\gamma$. Moreover, the parallel isomorphism depends only on the homotopy class of γ .

For $z_1 > z_2 > z_1 - z_2 > 0$ and each triple of V -modules W_1, W_2, W_3 , there is an associativity isomorphism:

$$\mathcal{A}_{P(z_1), P(z_2)}^{P(z_1-z_2), P(z_2)} : W_1 \boxtimes_{P(z_1)} (W_2 \boxtimes_{P(z_2)} W_3) \rightarrow (W_1 \boxtimes_{P(z_1-z_2)} W_2) \boxtimes_{P(z_2)} W_3,$$

which is characterized by

$$\overline{\mathcal{A}_{P(z_1), P(z_2)}^{P(z_1-z_2), P(z_2)}}(w_{(1)} \boxtimes_{P(z_1)} (w_{(2)} \boxtimes_{P(z_2)} w_{(3)})) = (w_{(1)} \boxtimes_{P(z_1-z_2)} w_{(2)}) \boxtimes_{P(z_2)} w_{(3)} \quad (3.3)$$

for $w_{(i)} \in W_i, i = 1, 2, 3$. The associativity isomorphism $\mathcal{A}$ of the braided tensor category is characterized by the following commutative diagram:

$$\begin{array}{ccc} W_1 \boxtimes_{P(z_1)} (W_2 \boxtimes_{P(z_2)} W_3) & \xrightarrow{\mathcal{T}_{\gamma_1} \circ (\text{id}_{W_1} \boxtimes_{P(z_1)} \mathcal{T}_{\gamma_2})} & W_1 \boxtimes (W_2 \boxtimes W_3) \\ \mathcal{A}_{P(z_1), P(z_2)}^{P(z_1-z_2), P(z_2)} \downarrow & & \downarrow \mathcal{A} \\ (W_1 \boxtimes_{P(z_1-z_2)} W_2) \boxtimes_{P(z_2)} W_3 & \xrightarrow{\mathcal{T}_{\gamma_2} \circ (\mathcal{T}_{\gamma_3} \boxtimes_{P(z_2)} \text{id}_{W_3})} & (W_1 \boxtimes W_2) \boxtimes W_3, \end{array} \quad (3.4)$$

where $\gamma_1, \gamma_2, \gamma_3$ are paths in $\mathbb{R}_+$ from $z_1, z_2, z_1 - z_2$ to 1, respectively.

There is also a braiding isomorphism, for $z > 0$, $\mathcal{R}_+^{P(z)} : W_1 \boxtimes_{P(z)} W_2 \rightarrow W_2 \boxtimes_{P(z)} W_1$ for each pair of V -modules W_1, W_2 , defined as

$$\overline{\mathcal{R}_+^{P(z)}}(w_1 \boxtimes_{P(z)} w_2) = e^{zL(-1)} \overline{\mathcal{T}_{\gamma_+}}(w_2 \boxtimes_{P(-z)} w_1), \quad (3.5)$$

where γ_+ is a path from $-z$ to z given by $\gamma_+(t) = -e^{i\pi t}z, t \in [0, 1]$. The inverse of $\mathcal{R}_+^{P(z)}$ is denoted by $\mathcal{R}_-^{P(z)}$, which is characterized by

$$\overline{\mathcal{R}_-^{P(z)}}(w_2 \boxtimes_{P(z)} w_1) = e^{zL(-1)} \overline{\mathcal{T}_{\gamma_-}}(w_1 \boxtimes_{P(-z)} w_2), \quad (3.6)$$

where γ_- is a path given by $\gamma_-(t) = -e^{-i\pi t}z$. We denote $\mathcal{R}_\pm^{P(1)}$ simply as $\mathcal{R}_\pm$. Accordingly, we define the twist by $\theta_W = e^{-2\pi i L(0)}$ for any V -module W .

For $z_1, z_2 \in \mathbb{R}_+$, the naturalness of $\mathcal{T}$ implies the commutativity of the following diagram:

$$\begin{array}{ccc} W_1 \boxtimes_{P(z_1)} W_2 & \xrightarrow{\mathcal{T}_\gamma} & W_1 \boxtimes_{P(z_2)} W_2 \\ \mathcal{R}_\pm^{P(z_1)} \downarrow & & \downarrow \mathcal{R}_\pm^{P(z_2)} \\ W_2 \boxtimes_{P(z_1)} W_1 & \xrightarrow{\mathcal{T}_\gamma} & W_2 \boxtimes_{P(z_2)} W_1, \end{array} \quad (3.7)$$

where γ is a path in $\mathbb{R}_+$ from z_1 to z_2 .

Let $\mathcal{V}_{W_1 W_2}^{W_3}$ denotes the space of intertwining operators of type $\binom{W_3}{W_1 W_2}$. For the chosen branch cut (1.16), the space $\mathcal{V}_{W_1 W_2}^{W_3}$ is canonically isomorphic to the space of intertwining maps denoted as $\mathcal{M}[P(z)]_{W_1 W_2}^{W_3}$. By the universal property of tensor product, the space $\mathcal{M}[P(z)]_{W_1 W_2}^{W_3}$ is canonically identified with the space $\text{Hom}_V(W_1 \boxtimes_{P(z)} W_2, W_3)$. Let $\mathcal{Y} \in \mathcal{V}_{W_1 W_2}^{W_3}$. We denote the corresponding morphism in $\text{Hom}_V(W_1 \boxtimes W_2, W_3)$ as $m_{\mathcal{Y}}^{P(z)}$. For $r \in \mathbb{Z}$, $\Omega_r : \mathcal{V}_{W_1 W_2}^{W_3} \rightarrow \mathcal{V}_{W_2 W_1}^{W_3}$ is an isomorphism defined as follows:

$$\Omega_r(\mathcal{Y})(w_2, x)w_1 = e^{xL(-1)}\mathcal{Y}(w_1, e^{(2r+1)\pi i}x)w_2$$

for all $w_1 \in W_1, w_2 \in W_2$. Then by Proposition 3.1 in [K2], where the definition of braiding is opposite to our choice here (recall (3.5)(3.6)), we obtain the following identities:

$$m_{\mathcal{Y}}^{P(z)} = m_{\Omega_0(\mathcal{Y})}^{P(z)} \circ \mathcal{R}_+^{P(z)} = m_{\Omega_{-1}(\mathcal{Y})}^{P(z)} \circ \mathcal{R}_-^{P(z)}. \quad (3.8)$$

The tensor product $V \otimes V$ is also a vertex operator algebra [FHL]. Moreover, it was shown in [HKo2] that $V \otimes V$ also satisfies the conditions in Theorem 0.1. Therefore, $\mathcal{C}_{V \otimes V}$, the category of $V \otimes V$ -modules, also has a structure of semisimple braided tensor category. For $z, \zeta \in \mathbb{C}^\times$, let $\boxtimes_{P(z)P(\zeta)}$ be the tensor product bifunctor in $\mathcal{C}_{V \otimes V}$ defined by

$$(A \otimes B) \boxtimes_{P(z)P(\zeta)} (C \otimes D) = (A \boxtimes_{P(z)} B) \otimes (C \boxtimes_{P(\zeta)} D),$$

where A, B, C, D are V -modules. We denote the bifunctor $\boxtimes_{P(1)P(1)}$ simply as $\boxtimes$.

There are a few different braiding structures on $\mathcal{C}_{V \otimes V}$ [K2]. We choose the one given by $\mathcal{R}_+ \otimes \mathcal{R}_-$. The twist $\theta_A : A \rightarrow A$, for $A \in \mathcal{C}_{V \otimes V}$, is given as follows

$$\theta_A = e^{-2\pi i L^L(0)} \otimes e^{2\pi i L^R(0)}. \quad (3.9)$$

An object A is said to have a trivial twist if $\theta_A = \text{id}_A$.

Now we recall the definition of (commutative) associative algebra in a braided tensor category $\mathcal{C}$ with tensor product $\otimes$, unit object $1_{\mathcal{C}}$, left unit isomorphism l_W and right unit isomorphism r_W for any object W , the associativity $\mathcal{A}$ and the braiding $\mathcal{R}$.

Definition 3.2. An associative algebra in $\mathcal{C}$ (or associative $\mathcal{C}$ -algebra) is an object A in $\mathcal{C}$ together with $\mu_A : A \otimes A \rightarrow A$ and a monomorphism $\iota_A : 1_{\mathcal{C}} \rightarrow A$ satisfying

1. *Associativity:* $\mu_A \circ (\mu_A \otimes \text{id}_A) \circ \mathcal{A} = \mu_A \circ (\text{id}_A \otimes \mu_A)$.
2. *Unit properties:* $\mu_A \circ (\iota_A \otimes \text{id}_A) \circ l_A^{-1} = \mu_A \circ (\iota_A \otimes \text{id}_A) \circ r_A^{-1} = \text{id}_A$.

A is called *commutative* if $\mu_A = \mu_A \circ \mathcal{R}$.

The following Theorem is proved in [HKo1].

Theorem 3.3. *The category of open-string vertex operator algebras over V is isomorphic to the category of associative $\mathcal{C}_V$ -algebras.*

The following Theorem is proved in [K2].

Theorem 3.4. *The category of conformal full field algebras over $V \otimes V$ is isomorphic to the category of commutative associative algebras in $\mathcal{C}_{V \otimes V}$ with a trivial twist.*

We are interested in studying the relation between above two algebras as ingredients of an open-closed field algebra over V . Notice that these two algebras live in different categories. So we will first discuss a functor between these two categories.

Recall a functor $T_{P(z)} : \mathcal{C}_{V \otimes V} \rightarrow \mathcal{C}_V$ [HL4]. In particular, for W_1, W_2 being V -module, $T_{P(z)}(W_1 \otimes W_2) = W_1 \boxtimes_{P(z)} W_2$. We will simply write $T_{P(1)}$ as T . Let

$$\varphi_0 := l_V^{-1} = r_V^{-1} : V \rightarrow V \boxtimes V = T(V \otimes V). \quad (3.10)$$

For each four V -modules $W_i^L, W_i^R, i = 1, 2$, notice that

$$T(W_1^L \otimes W_1^R) \boxtimes T(W_2^L \otimes W_2^R) = (W_1^L \boxtimes W_1^R) \boxtimes (W_2^L \boxtimes W_2^R), \quad (3.11)$$

$$T((W_1^L \otimes W_2^R) \boxtimes (W_2^L \otimes W_2^R)) = (W_1^L \boxtimes W_2^L) \boxtimes (W_1^R \boxtimes W_2^R). \quad (3.12)$$


We define $\varphi_2 : (W_1^L \boxtimes W_1^R) \boxtimes (W_2^L \boxtimes W_2^R) \rightarrow (W_1^L \boxtimes W_2^L) \boxtimes (W_1^R \boxtimes W_2^R)$ by

$$\varphi_2 := \mathcal{A} \circ (\text{id}_{W_1} \boxtimes \mathcal{A}^{-1}) \circ (\text{id}_{W_1} \boxtimes \mathcal{R}_- \boxtimes \text{id}_{W_4}) \circ (\text{id}_{W_1} \boxtimes \mathcal{A}) \circ \mathcal{A}^{-1}. \quad (3.13)$$

The above definition of φ_2 can be naturally extended to a morphism $T(A) \boxtimes T(B) \rightarrow T(A \boxtimes B)$ for each pair of objects A and B in $\mathcal{C}_{V \otimes V}$. We still denote the extended morphism as φ_2 . The following result is clear.

Lemma 3.5. *T together with φ_0, φ_2 given in (3.10) (3.13) is a monoidal functor.*

For any four objects $W_i, i = 1, 2, 3, 4$ in $\mathcal{C}_V$, we define a morphism σ in $\text{Hom}((W_1 \boxtimes W_2) \boxtimes (W_3 \boxtimes W_4), (W_3 \boxtimes W_4) \boxtimes (W_1 \boxtimes W_2))$ according to the following graph:



(3.14)

Clearly, for any $A \in \mathcal{C}_{V \otimes V}$, σ can be extended to an automorphism on $T(A) \boxtimes T(A)$, denoted as σ_A .

Proposition 3.6. *Let (A, μ_A, ι_A) be a commutative algebra in $\mathcal{C}_{V \otimes V}$. Let $\mu_{T(A)} := T(\mu_A) \circ \varphi_2$ and $\iota_{T(A)} := T(\iota_A) \circ \varphi_0$. Then $(T(A), \mu_{T(A)}, \iota_{T(A)})$ is an associative $\mathcal{C}_V$ -algebra satisfying the following commutativity:*

$$\mu_{T(A)} = \mu_{T(A)} \circ \sigma_A. \quad (3.15)$$

Proof. That $(T(A), \mu_{T(A)}, \iota_{T(A)})$ is an associative $\mathcal{C}_V$ -algebra follows from Lemma 3.5. We only prove (3.15). Let us assume $A = W_1 \otimes W_2$ with W_1, W_2 being V -modules (the proof for general A is the same). Consider the following diagram:


$$\begin{array}{ccccc}
(W_1 \boxtimes W_2) \boxtimes (W_1 \boxtimes W_2) & \xrightarrow{\varphi_2} & (W_1 \boxtimes W_1) \boxtimes (W_2 \boxtimes W_2) & \xrightarrow{T(\mu_{cl})} & W_1 \boxtimes W_2 \\
\downarrow \sigma & & \downarrow \mathcal{R}_+ \boxtimes \mathcal{R}_- & & \downarrow \text{id} \\
(W_1 \boxtimes W_2) \boxtimes (W_1 \boxtimes W_2) & \xrightarrow{\varphi_2} & (W_1 \boxtimes W_1) \boxtimes (W_2 \boxtimes W_2) & \xrightarrow{T(\mu_{cl})} & W_1 \boxtimes W_2 .
\end{array} \quad (3.16)$$

The left subdiagram is commutative because two paths corresponding to the same braiding; the right subdiagram is commutative because of the commutativity of A . Hence the above diagram is commutative. (3.15) follows from the commutativity of (3.16). ■


3.2 Open-closed $\mathcal{C}_V | \mathcal{C}_{V \otimes V}$ -algebras

Let $((V_{cl}, m_{cl}, \iota_{cl}), (V_{op}, Y_{op}, \iota_{op}), \mathbb{Y}_{cl-op})$ be an open-closed field algebra over V throughout this subsection.

By Theorem 3.4, $(V_{cl}, m_{cl}, \iota_{cl})$, a conformal full field algebra over $V \otimes V$ is equivalent to a commutative associative algebra with a trivial twist in $\mathcal{C}_{V \otimes V}$ with braiding $\mathcal{R}_{+-}$. We denoted this $\mathcal{C}_{V \otimes V}$ -algebra as a triple $(V_{cl}, \mu_{cl}, \iota_{cl})$, where $\mu_{cl} = m_{\mathbb{Y}_f}$ (recall (1.31)). By Proposition 3.6, $(T(V_{cl}), \nu_{T(V_{cl})}, \iota_{T(V_{cl})})$ is an associative $\mathcal{C}_V$ -algebra. The property of algebra $T(V_{cl})$ can be expressed in the following graphic equations:




$$\text{Diagrammatic equation (3.17)} \quad (3.17)$$



$$\text{Diagrammatic equation (3.18)} \quad (3.18)$$

By Theorem 3.3, $(V_{op}, Y_{op}, \iota_{op})$, an open-string vertex operator algebra over V is equivalent to an associative $\mathcal{C}_V$ -algebra, denoted as a triple $(V_{op}, \mu_{op}, \iota_{op})$, where $\mu_{op} = m_{Y_{op}^f}$ (recall (1.14) and (1.15)). The defining properties of $(V_{op}, \mu_{op}, \iota_{op})$ can be expressed in the following graphic equations:



$$\text{Diagrammatic equation (3.19)} \quad (3.19)$$

It remains to study the categorical formulation of the only remaining data $\mathbb{Y}_{cl-op}$ and its properties.

By the chirality splitting properties and the associativity of intertwining operator algebra [H7], there exist intertwining operators $\mathcal{Y}^{(5)}, \mathcal{Y}^{(6)}$ such that

$$\langle v', \mathbb{Y}_{cl-op}(u \otimes \bar{u}, z, \zeta)v \rangle = \langle v', \mathcal{Y}^{(5)}(\mathcal{Y}^{(6)}(u^L, z - \zeta)u^R, \zeta)v \rangle \quad (3.20)$$

for $v' \in V'_{op}, v \in V_{op}, u^L \otimes u^R \in V_{cl}$ and $\zeta > z - \zeta > 0$. For $u^L \otimes u^R \in V_{cl}$, we have $u^L \boxtimes_{P(z-\zeta)} u^R \in \overline{T_{P(z-\zeta)}(V_{cl})}$.

We define a map $\mu_{cl-op}^{P(z-\zeta)P(\zeta)} : T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} V_{op} \rightarrow V_{op}$ as follows:

$$\mu_{cl-op}^{P(z-\zeta)P(\zeta)} := m_{\mathcal{Y}^{(5)}}^{P(\zeta)} \circ (m_{\mathcal{Y}^{(6)}}^{P(z-\zeta)} \boxtimes_{P(\zeta)} \text{id}_{V_{op}}). \quad (3.21)$$

By the convergence and analytic properties of the right hand side of (3.20) and the fact that module maps are continuous with respect to the topology on graded vector spaces defined in Section 1, we obtain, for $z > \zeta > z - \zeta > 0$, $u^L \otimes u^R \in V_{cl}, v \in V_{op}$, the following identity:

$$\overline{\mu_{cl-op}^{P(z-\zeta)P(\zeta)}}((u^L \boxtimes_{P(z-\zeta)} u^R) \boxtimes_{P(\zeta)} v) = \mathcal{Y}^{(5)}(\mathcal{Y}^{(6)}(u^L, z - \zeta)u^R, \zeta)v, \quad (3.22)$$

where $\overline{\mu_{cl-op}^{P(z-\zeta)P(\zeta)}}$ is the unique extension of $\mu_{cl-op}^{P(z-\zeta)P(\zeta)}$ to the algebraic completion of $T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} V_{op}$. Moreover, by Proposition 1.2 in [H9] (or the nondegeneracy of intertwining operator algebra [K1]), such V -module map is unique.

Let γ_1 be a path in $\mathbb{R}_+$ from 1 to $z - \zeta$ and γ_2 a path in $\mathbb{R}_+$ from 1 to ζ . Then we define a map $\mu_{cl-op} : T(V_{cl}) \boxtimes V_{op} \rightarrow V_{op}$ as follows:

$$\mu_{cl-op} := \mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ \mathcal{T}_{\gamma_2} \circ (\mathcal{T}_{\gamma_1} \boxtimes \text{id}_{V_{op}}). \quad (3.23)$$

Since $\mathcal{T}_\gamma$ depends on path only homotopically, it is clear that above definition of μ_{cl-op} is independent of z, ζ in $\mathbb{R}_+$ and paths γ_1, γ_2 in $\mathbb{R}_+$. In particular, one can choose γ_1 and γ_2 to be the straight line between 1 and $z - \zeta, \zeta$ respectively.


By Theorem 1.28, to obtain a categorical formulation of open-closed field algebra over V is enough to study the categorical formulations of the unit property (1.82), Associativity I (1.52), Associativity II (1.56) and Commutativity I in Proposition 1.18.

We first consider the property (1.82).

Proposition 3.7. *The condition (1.82) is equivalent to the following condition:*

$$\mu_{cl-op} \circ ((T(\iota_{V_{cl}}) \circ \varphi_0) \boxtimes \text{id}_{V_{op}}) \circ l_{V_{op}}^{-1} = \text{id}_{V_{op}}, \quad (3.24)$$

which can also be expressed by the following graphic equation:



$$\text{Diagram} = \text{Diagram} \quad (3.25)$$

Proof. First, (1.82) is equivalent to the following condition:

$$\mathbb{Y}_{cl-op}(\mathbf{1}_{cl}; z, \zeta) = \text{id}_{V_{op}}, \quad \text{for } z > \zeta > 0. \quad (3.26)$$

Recall that $\mathbf{1}_{cl} = \iota_{V_{cl}}(\mathbf{1} \otimes \mathbf{1})$. Replacing $u^L \otimes u^R$ by $\mathbf{1}_{cl}$ in (3.20), one can see that both sides of equation (3.20) are independent of z and ζ . Hence (3.26) holds for all $z, \zeta \in \mathbb{R}_+$. Using (3.20) and (3.22), we obtain

$$\overline{\mu_{cl-op}^{P(z-\zeta)P(\zeta)}}(T_{P(z-\zeta)}(\mathbf{1}_{cl}) \boxtimes_{P(\zeta)} v) = v \quad (3.27)$$

for $v \in V_{op}$ and $z, \zeta \in \mathbb{R}_+$. Therefore, we have, for $v \in V_{op}$,

$$\begin{aligned} \mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ T_{P(z-\zeta)}(\iota_{V_{cl}}) \boxtimes_{P(\zeta)} \text{id}_{V_{op}}((\mathbf{1} \boxtimes_{P(z-\zeta)} \mathbf{1}) \boxtimes_{P(\zeta)} v) \\ = \mu_{cl-op}^{P(z-\zeta)P(\zeta)}(T_{P(z-\zeta)}(\mathbf{1}_{cl}) \boxtimes_{P(\zeta)} v) \\ = v, \end{aligned} \quad (3.28)$$

which can be further expressed equivalently as

$$\begin{aligned} \text{id}_{V_{op}} &= \mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ ((T_{P(z-\zeta)}(\iota_{V_{cl}}) \circ \mathcal{T}_{\gamma_1}) \boxtimes_{P(\zeta)} \text{id}_{V_{op}}) \circ \mathcal{T}_{\gamma_2} \circ (l_V^{-1} \boxtimes \text{id}_{V_{op}}) \circ l_{V_{op}}^{-1} \\ &= \mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ (\mathcal{T}_{\gamma_1} \boxtimes_{P(\zeta)} \text{id}_{V_{op}}) \circ \mathcal{T}_{\gamma_2} \circ ((T(\iota_{V_{cl}}) \circ \varphi_0) \boxtimes \text{id}_{V_{op}}) \circ l_{V_{op}}^{-1} \\ &= \mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ \mathcal{T}_{\gamma_2} \circ (\mathcal{T}_{\gamma_1} \boxtimes \text{id}_{V_{op}}) \circ ((T(\iota_{V_{cl}}) \circ \varphi_0) \boxtimes \text{id}_{V_{op}}) \circ l_{V_{op}}^{-1} \\ &= \mu_{cl-op} \circ ((T(\iota_{V_{cl}}) \circ \varphi_0) \boxtimes \text{id}_{V_{op}}) \circ l_{V_{op}}^{-1} \end{aligned} \quad (3.29)$$

where γ_1 and γ_2 are paths in $\mathbb{R}_+$ from 1 to $z - \zeta$ and ζ respectively.


Conversely, from (3.27), (3.28) and (3.29), it is clear that (3.24) or (3.25) also implies (1.82). $\blacksquare$

Now we consider the associativity II (recall Proposition 1.17).

Proposition 3.8. *The associativity II is equivalent to the following condition:*

$$\mu_{cl-op} \circ (\text{id}_{T(V_{cl})} \boxtimes \mu_{cl-op}) = \mu_{cl-op} \circ ((T(\mu_{cl}) \circ \varphi_2) \boxtimes \text{id}_{V_{op}}) \circ \mathcal{A}, \quad (3.30)$$

which can also be expressed by the following graphic equation



$$(3.31)$$

Proof. Using the convergence property of products and iterates of intertwining operators of V , it is not hard to show that, in the domain

$$D_3 := \{(z_1, \zeta_1, z_2, \zeta_2) | z_1 > \zeta_1 > z_2 > \zeta_2 > 0, 2\zeta_2 > z_2, 2\zeta_1 > z_1 + z_2\}, \quad (3.32)$$

we have

$$\begin{aligned} Y_{cl-op}(u_1^L \otimes u_1^R; z_1, \zeta_1) Y_{cl-op}(u_1^L \otimes u_1^R; z_1, \zeta_1) v \\ = \mathcal{Y}^{(5)}(\mathcal{Y}^{(6)}(u_1^L, z_1 - \zeta_1) u_1^R, \zeta_1) \mathcal{Y}^{(5)}(\mathcal{Y}^{(6)}(u_2^L, z_2 - \zeta_2) u_2^R, \zeta_2) v \end{aligned} \quad (3.33)$$

for $u_1^L \otimes u_1^R, u_2^L \otimes u_2^R \in V_{cl}$ and $v \in V_{op}$.

By (3.21)(3.22) and the fact that module maps are continuous, we obtain

$$\begin{aligned} Y_{cl-op}(u_1^L \otimes u_1^R; z_1, \zeta_1) Y_{cl-op}(u_1^L \otimes u_1^R; z_1, \zeta_1) v \\ = \overline{\mu_{cl-op}^{P(z_1-\zeta_1)P(\zeta_1)} \circ (\text{id}_{T_{P(z_1-\zeta_1)}(V_{cl})} \boxtimes_{P(\zeta_1)} \mu_{cl-op}^{P(z_2-\zeta_2)P(\zeta_2)})} (\\ (u_1^L \boxtimes_{P(z_1-\zeta_1)} u_1^R) \boxtimes_{P(\zeta_1)} ((u_2^L \boxtimes_{P(z_2-\zeta_2)} u_2^R) \boxtimes_{P(\zeta_2)} v)) \end{aligned} \quad (3.34)$$

for $(z_1, \zeta_1, z_2, \zeta_2)$ in the domain D_3 . Moreover, by the universal property of tensor product and the nondegeneracy of intertwining operator algebra [K1],

$$\mu_{cl-op}^{P(z_1-\zeta_1)P(\zeta_1)} \circ (\text{id}_{T_{P(z_1-\zeta_1)}(V_{cl})} \boxtimes_{P(\zeta_1)} \mu_{cl-op}^{P(z_2-\zeta_2)P(\zeta_2)}) \quad (3.35)$$

is the unique morphism in $\text{Hom}(T_{P(z_1-\zeta_1)}(V_{cl}) \boxtimes_{P(\zeta_1)} (T_{P(z_2-\zeta_2)}(V_{cl}) \boxtimes_{P(\zeta_2)} V_{op}), V_{op})$ such that the identity (3.34) holds in D_3 .

On the other hand of (1.56), it is proved in [HKO2] that $\mathbb{Y}$ can be expanded as follows:

$$\mathbb{Y}_{an}(u_1^L \otimes u_1^R; z, \zeta) u_2^L \otimes u_2^R = \sum_{i=1}^N \mathcal{Y}_i^L(u_1^L, z) u_2^L \otimes \mathcal{Y}_i^R(u_1^R, \zeta) u_2^R \quad (3.36)$$

for some $N \in \mathbb{Z}_+$. There is a unique morphism

$$\mu_{cl}^{P(z_1-z_2)P(\zeta_1-\zeta_2)} \in \text{Hom}_{\mathcal{C}_{V \otimes V}}(V_{cl} \boxtimes_{P(z_1-z_2)P(\zeta_1-\zeta_2)} V_{cl}, V_{cl})$$

such that, for $u, v \in V_{cl}$,

$$\overline{\mu_{cl}^{P(z_1-z_2)P(\zeta_1-\zeta_2)}}(u \boxtimes_{P(z_1-z_2)P(\zeta_1-\zeta_2)} v) = \mathbb{Y}_{an}(u; z_1 - z_2, \zeta_1 - \zeta_2)v. \quad (3.37)$$

By the convergence property of intertwining operator algebra, it not hard to see that, in the domain

$$D_4 := \{(z_1, \zeta_1, z_2, \zeta_2) | z_1 > z_2 > \zeta_1 > \zeta_2 > 0, 2\zeta_2 > z_1, 2z_2 > z_1 + \zeta_1\}, \quad (3.38)$$

we have

$$\begin{aligned} & Y_{cl-op}(\mathbb{Y}_{an}(u_1^L \otimes u_1^R; z_1 - z_2, \zeta_1 - \zeta_2) u_2^L \otimes u_2^R; z_2, \zeta_2)v \\ &= \sum_{i=1}^N Y_{cl-op}(\mathcal{Y}_i^L(u_1^L, z_1 - z_2) u_2^L \otimes \mathcal{Y}_i^R(u_1^R, \zeta_1 - \zeta_2) u_2^R; z_2, \zeta_2)v \\ &= \sum_{i=1}^N \mathcal{Y}^{(5)}(\mathcal{Y}^{(6)}(\mathcal{Y}_i^L(u_1^L, z_1 - z_2) u_2^L, z_2 - \zeta_2) \mathcal{Y}_i^R(u_1^R, \zeta_1 - \zeta_2) u_2^R, \zeta_2)v. \end{aligned} \quad (3.39)$$

Combining (3.39) with (3.37), (3.20) and (3.22) and the fact that module maps are continuous, we obtain the following identity:

$$\begin{aligned} & \mathbb{Y}_{cl-op}(\mathbb{Y}_{an}(u_1^L \otimes u_1^R; z_1 - z_2, \zeta_1 - \zeta_2) u_2^L \otimes u_2^R; z_2, \zeta_2)v \\ &= \overline{\mu_{cl-op}^{P(z_2-\zeta_2)P(\zeta_2)} \circ (T_{P(z_2-\zeta_2)}(\mu_{cl}^{P(z_1-z_2)P(\zeta_1-\zeta_2)}) \boxtimes_{P(\zeta_2)} \text{id}_{V_{op}})}(\\ & \quad ((u_1^L \boxtimes_{P(z_1-z_2)} u_2^L) \boxtimes_{P(z_2-\zeta_2)} (u_1^R \boxtimes_{P(\zeta_1-\zeta_2)} u_2^R)) \boxtimes_{P(\zeta_2)} v) \end{aligned} \quad (3.40)$$

for $(z_1, \zeta_1, z_2, \zeta_2)$ in the domain D_4 . Moreover, by the universal property of tensor product and the nondegeneracy of intertwining operator algebra [K1],

$$\mu_{cl-op}^{P(z_2-\zeta_2)P(\zeta_2)} \circ (T_{P(z_2-\zeta_2)}(\mu_{cl}^{P(z_1-z_2)P(\zeta_1-\zeta_2)}) \boxtimes_{P(\zeta_2)} \text{id}_{V_{op}}) \quad (3.41)$$

is the unique morphism in $\text{Hom}(T_{P(z_2-\zeta_2)}(V_{cl} \boxtimes_{P(z_1-z_2)P(\zeta_1-\zeta_2)} V_{cl}) \boxtimes_{P(\zeta_2)} V_{op}, V_{op})$ such that the identity (3.40) holds in D_4 .

Notice that the domains D_3 and D_4 are disjoint. Now we fix a point $(z_1, \zeta_1, z_2, \zeta_2)$ in the domain

$$\{(z_1, \zeta_1, z_2, \zeta_2) | z_1 > \zeta_1 > z_2 > \zeta_2 > 0, 2\zeta_2 > z_1, 2\zeta_1 > z_1 + z_2, 2z_2 > \zeta_1 + \zeta_2\}, \quad (3.42)$$

which is a subdomain of D_3 . Let $\tilde{z}_2 = \zeta_1, \tilde{\zeta}_1 = z_2$. Then the quadruple $(z_1, \tilde{\zeta}_1, \tilde{z}_2, \zeta_2)$ is in the domain D_4 .

By the analytic properties guaranteed by Theorem 1.28, both sides of associativity (1.56) can be uniquely extended to the boundary $(z_1, \zeta_1, z_2, \zeta_2) \in \mathbb{R}_+^4 \cap M_{\mathbb{C}^\times}^4$. Moreover,

$$\mathbb{Y}_{cl-op}(u_1^L \otimes u_1^R; z_1, \zeta_1) \mathbb{Y}_{cl-op}(u_2^L \otimes u_2^R; z_2, \zeta_2) v,$$

and

$$\mathbb{Y}_{cl-op}(\mathbb{Y}_{an}(u_1^L \otimes u_1^R; z_1 - \tilde{z}_2, \tilde{\zeta}_1 - \zeta_2) u_2^L \otimes u_2^R; \tilde{z}_2, \zeta_2) v,$$

can be obtained from each other by analytic continuation along the following path

Meanwhile, if we start from the element

$$(u_1^L \boxtimes_{P(z_1-\zeta_1)} u_1^R) \boxtimes_{P(\zeta_1)} ((u_2^L \boxtimes_{P(z_2-\zeta_2)} u_2^R) \boxtimes_{P(\zeta_2)} v)$$

in $\overline{T_{P(z_1-\zeta_1)}(V_{cl}) \boxtimes_{P(\zeta_1)} (T_{P(z_2-\zeta_2)}(V_{cl}) \boxtimes_{P(\zeta_2)} V_{op})}$ and apply associativity isomorphisms repeatedly and braiding isomorphism on it, we obtain

$$\begin{aligned} & \xrightarrow{(\mathcal{A}_{P(z_1)P(\zeta_1)}^{P(z_1-\zeta_1)P(\zeta_1)})^{-1}} u_1^L \boxtimes_{P(z_1)} (u_1^R \boxtimes_{P(\zeta_1)} ((u_2^L \boxtimes_{P(z_2-\zeta_2)} u_2^R) \boxtimes_{P(\zeta_2)} v)), \\ & \xrightarrow{\mathcal{A}_{P(\zeta_1)P(\zeta_2)}^{P(\zeta_1-\zeta_2)P(\zeta_2)}} u_1^L \boxtimes_{P(z_1)} ((u_1^R \boxtimes_{P(\zeta_1-\zeta_2)} (u_2^L \boxtimes_{P(z_2-\zeta_2)} u_2^R)) \boxtimes_{P(\zeta_2)} v), \\ & \xrightarrow{\mathcal{A}_{P(\zeta_1-\zeta_2)P(z_2-\zeta_2)}^{P(\zeta_1-z_2)P(z_2-\zeta_2)}} u_1^L \boxtimes_{P(z_1)} (((u_1^R \boxtimes_{P(\zeta_1-z_2)} u_2^L) \boxtimes_{P(z_2-\zeta_2)} u_2^R) \boxtimes_{P(\zeta_2)} v), \\ & \xrightarrow{\mathcal{R}_-^{P(\zeta_1-z_2)}} u_1^L \boxtimes_{P(z_1)} ((e^{(\zeta_1-z_2)L(-1)} \cdot \overline{T_{\gamma_-}}(u_2^L \boxtimes_{P(-\zeta_1+z_2)} u_1^R) \boxtimes_{P(z_2-\zeta_2)} u_2^R) \boxtimes_{P(\zeta_2)} v), \end{aligned} \quad (3.44)$$

where we have ignored the obvious identity maps and γ_- is a path from $-\zeta_1 + z_2$ to $\zeta_1 - z_2$ given by $\gamma_-(t) = (-\zeta_1 + z_2)e^{-\pi it}$, $t \in [0, 1]$. Now we analytic continue the last line of (3.44) along the path (3.43), we obtain

$$u_1^L \boxtimes_{P(z_1)} (((u_2^L \boxtimes_{P(\zeta_1-z_2)} u_1^R) \boxtimes_{P(z_2-\zeta_2)} u_2^R) \boxtimes_{P(\zeta_2)} v), \quad (3.45)$$

which is nothing but the third line of (3.44) except u_1^R and u_2^L being exchanged. Note that the last line of (3.44) and (3.45) are elements in the algebraic completion of the

same V -module. We denote this V -module as W . Now we further apply associativity morphisms on (3.45) and use $z_2 = \tilde{\zeta}_1$ and $\zeta_1 = \tilde{z}_2$. We then obtain

$$\begin{aligned} & \overline{(\mathcal{A}_{P(\tilde{z}_2-\tilde{\zeta}_2)}^{P(\tilde{z}_2-\tilde{\zeta}_2)P(\tilde{\zeta}_1-\zeta_2)})^{-1}} \rightarrow u_1^L \boxtimes_{P(z_1)} ((u_2^L \boxtimes_{P(\tilde{z}_2-\zeta_2)} (u_1^R \boxtimes_{P(\tilde{\zeta}_1-\zeta_2)} u_2^R)) \boxtimes_{P(\zeta_2)} v), \\ & \overline{\mathcal{A}_{P(z_1)P(\zeta_2)}^{P(z_1-\zeta_2)P(\zeta_2)}} \rightarrow (u_1^L \boxtimes_{P(z_1-\zeta_2)} (u_2^L \boxtimes_{P(\tilde{z}_2-\zeta_2)} (u_1^R \boxtimes_{P(\tilde{\zeta}_1-\zeta_2)} u_2^R))) \boxtimes_{P(\zeta_2)} v, \\ & \overline{\mathcal{A}_{P(z_1-\zeta_2)P(\tilde{z}_2-\zeta_2)}^{P(z_1-\tilde{z}_2)P(\tilde{z}_2-\zeta_2)}} \rightarrow ((u_1^L \boxtimes_{P(z_1-\tilde{z}_2)} u_2^L) \boxtimes_{P(\tilde{z}_2-\zeta_2)} (u_1^R \boxtimes_{P(\tilde{\zeta}_1-\zeta_2)} u_2^R)) \boxtimes_{P(\zeta_2)} v. \end{aligned} \quad (3.46)$$

Let m be the morphism $W \rightarrow V_{op}$ such that

$$\begin{aligned} (3.35) &= m \circ \mathcal{R}_-^{P(\zeta_1-z_2)} \circ \mathcal{A}_{P(\zeta_1-\zeta_2)P(z_2-\zeta_2)}^{P(\zeta_1-z_2)P(z_2-\zeta_2)} \circ \\ & \quad \mathcal{A}_{P(\zeta_1)P(\zeta_2)}^{P(\zeta_1-\zeta_2)P(\zeta_2)} \circ (\mathcal{A}_{P(z_1)P(\zeta_1)}^{P(z_1-\zeta_1)P(\zeta_1)})^{-1}. \end{aligned} \quad (3.47)$$

If we apply $\overline{m}$ on both the last line of (3.44) and (3.45), the two images of $\overline{m}$ are clearly the analytic continuation of each other along the path (3.43). On the other hand, combining this fact with (3.40) and (3.46), we obtain that

$$m = (3.41) \circ \mathcal{A}_{P(z_1-\zeta_2)P(\tilde{z}_2-\zeta_2)}^{P(z_1-\tilde{z}_2)P(\tilde{z}_2-\zeta_2)} \circ \mathcal{A}_{P(z_1)P(\zeta_2)}^{P(z_1-\zeta_2)P(\zeta_2)} \circ (\mathcal{A}_{P(\tilde{z}_2-\zeta_2)}^{P(\tilde{z}_2-\tilde{\zeta}_2)P(\tilde{\zeta}_1-\zeta_2)})^{-1} \quad (3.48)$$

because the extensions of both sides of (3.48), applied on (3.45), give the same element in $\overline{V_{op}}$. Therefore, we further obtain from (3.47) and (3.48) the following identity:

$$\begin{aligned} (3.35) &= (3.41) \circ \mathcal{A}_{P(z_1-\zeta_2)P(\tilde{z}_2-\zeta_2)}^{P(z_1-\tilde{z}_2)P(\tilde{z}_2-\zeta_2)} \circ \mathcal{A}_{P(z_1)P(\zeta_2)}^{P(z_1-\zeta_2)P(\zeta_2)} \circ (\mathcal{A}_{P(\tilde{z}_2-\zeta_2)}^{P(\tilde{z}_2-\tilde{\zeta}_2)P(\tilde{\zeta}_1-\zeta_2)})^{-1} \\ & \quad \circ \mathcal{R}_-^{P(\zeta_1-z_2)} \circ \mathcal{A}_{P(\zeta_1-\zeta_2)P(z_2-\zeta_2)}^{P(\zeta_1-z_2)P(z_2-\zeta_2)} \circ \mathcal{A}_{P(\zeta_1)P(\zeta_2)}^{P(\zeta_1-\zeta_2)P(\zeta_2)} \circ (\mathcal{A}_{P(z_1)P(\zeta_1)}^{P(z_1-\zeta_1)P(\zeta_1)})^{-1}. \end{aligned} \quad (3.49)$$

Using the commutative diagram (3.4)(3.7) and the definition of φ_2 (recall (3.13)), it is easy to see that (3.49) implies the commutativity of the following diagram:

$$\begin{array}{ccc} T_{P(z_1-\zeta_1)}(V_{cl}) \boxtimes_{P(\zeta_1)} (T_{P(z_2-\zeta_2)}(V_{cl}) \boxtimes_{P(\zeta_2)} V_{op}) & & (3.50) \\ \downarrow f_1 & \searrow (3.35) & \\ T(V_{cl}) \boxtimes (T(V_{cl}) \boxtimes V_{op}) & & \\ \downarrow \mathcal{A} & \searrow \text{dotted} & \\ (T(V_{cl}) \boxtimes T(V_{cl})) \boxtimes V_{op} & & V_{op} \\ \downarrow \varphi_2 \boxtimes \text{id}_{V_{op}} & \searrow \text{dotted} & \\ T(V_{cl} \boxtimes V_{cl}) \boxtimes V_{op} & & \\ \downarrow g_1 & \searrow (3.41)|_{z_2=\tilde{z}_2, \zeta_1=\tilde{\zeta}_1} & \\ T_{P(\tilde{z}_2-\zeta_2)}(V_{cl} \boxtimes_{P(z_1-\tilde{z}_2)P(\tilde{\zeta}_1-\zeta_2)} V_{cl}) \boxtimes_{P(\zeta_2)} V_{op} & & \end{array}$$

where

$$f_1 = \text{id}_{T(V_{cl})} \boxtimes (\mathcal{T}_{\gamma_4} \boxtimes \text{id}_{H_{op}}) \circ \mathcal{T}_{\gamma_2} \boxtimes \mathcal{T}_{\gamma_3} \circ \mathcal{T}_{\gamma_1} \quad (3.51)$$

$$g_1 = T_{P(z_2-\zeta_2)}(\mathcal{T}_{\gamma_7} \boxtimes \mathcal{T}_{\gamma_8}) \boxtimes_{P(\zeta_2)} \text{id}_{H_{op}} \circ (\mathcal{T}_{\gamma_6} \boxtimes_{P(\zeta_2)} \text{id}_{V_{op}}) \circ \mathcal{T}_{\gamma_5} \quad (3.52)$$

in which $\gamma_i, i = 1, \dots, 4$ are paths in $\mathbb{R}_+$ from $\zeta_1, z_1 - \zeta_1, \zeta_2, z_2 - \zeta_2$ to 1 respectively and $\gamma_i, i = 5, \dots, 8$ are paths in $\mathbb{R}_+$ from 1 to $\zeta_2, z_2 - \zeta_2, z_1 - z_2, \zeta_1 - \zeta_2$ respectively.

Using (3.23), it is easy to see that

$$\begin{aligned}\mu_{cl-op} \circ (\text{id}_{T(V_{cl})} \boxtimes \mu_{cl-op}) &= (3.35) \circ f_1^{-1}, \\ \mu_{cl-op} \circ (T(\mu_{cl}) \boxtimes \text{id}_{V_{op}}) &= (3.41) \circ g_1.\end{aligned}\tag{3.53}$$

(3.53) together with the commutative diagram (3.50) implies (3.30), which is nothing but the commutativity of the subdiagram in the middle of (3.50).


Conversely, (3.30) implies the commutativity of the diagram (3.50). It is easy to see that above arguments can be reversed. Therefore, we can also obtain the associativity II (1.56) from (3.30). $\blacksquare$

We now study the categorical formulations of the rest conditions needed in Theorem 1.28. The proof of them are essentially the same as that of Proposition 3.8. So we will only sketch the proofs below.

Proposition 3.9. *The associativity I (recall Proposition 1.16) is equivalent to the following condition:*

$$\mu_{cl-op}(\text{id}_{T(V_{cl})} \boxtimes \mu_{op}) = \mu_{op}(\mu_{cl-op} \boxtimes \text{id}_{V_{op}}) \circ \mathcal{A}\tag{3.54}$$

which can also be expressed in the following graph:


(3.55)

Proof. The left hand side of (1.52) gives arise to a morphism

$$\mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ (\text{id}_{T_{P(z-\zeta)}(V_{cl})} \boxtimes_{P(\zeta)} \mu_{op}^{P(r)})\tag{3.56}$$

in $\text{Hom}(T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} (V_{op} \boxtimes_{P(r)} V_{op}), V_{op})$. The right hand side of (1.52) gives arise to a morphism

$$\mu_{op}^{P(r)} \circ (\mu_{cl-op}^{P(z-\zeta)P(\zeta-r)} \boxtimes_{P(\zeta-r)} \text{id}_{V_{op}})\tag{3.57}$$

in $\text{Hom}((T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} V_{op}) \boxtimes_{P(r)} V_{op}, V_{op})$.

For z, ζ, r in a proper subdomain of $z > \zeta > r > 0$, using similar arguments as in Proposition 3.8, we obtain that the associativity (1.52), for some $z, \zeta, r, \tilde{r} \in \mathbb{R}_+$, implies the following commutative diagram:

$$\begin{array}{ccc} T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} (V_{op} \boxtimes_{P(r)} V_{op}) & & (3.58) \\ \downarrow f_2 & \searrow (3.56) & \\ T(V_{cl}) \boxtimes (V_{op} \boxtimes V_{op}) & \cdots \cdots \cdots \rightarrow & V_{op} \\ \downarrow \mathcal{A} & \nearrow (3.57) & \\ (T(V_{cl}) \boxtimes V_{op}) \boxtimes V_{op} & & \\ \downarrow g_2 & \nearrow & \\ (T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta-r)} V_{op}) \boxtimes_{P(r)} V_{op}, & & \end{array}$$

where

$$f_2 = (\mathcal{T}_{\gamma_2} \boxtimes \mathcal{T}_{\gamma_3}) \circ \mathcal{T}_{\gamma_1}, \quad (3.59)$$

$$g_2 = \mathcal{T}_{\gamma_6} \boxtimes_{P(\zeta)} \text{id}_{V_{op}} \boxtimes_{P(r)} \text{id}_{V_{op}} \circ (\mathcal{T}_{\gamma_5} \boxtimes_{P(r)} \text{id}_{V_{op}}) \circ \mathcal{T}_{\gamma_4}, \quad (3.60)$$

where $\gamma_i, i = 1, 2, 3$ are paths in $\mathbb{R}_+$ from $\zeta, z - \zeta$ and r to 1 respectively and $\gamma_i, i = 4, 5, 6$ are paths in $\mathbb{R}_+$ from 1 to $r, \zeta - r$ and $z - \zeta$ respectively. The commutativity of outside loop in (3.58) implies immediately the commutativity of the subdiagram in the middle of (3.58), which is nothing but the identity (3.54) or (3.55).

Conversely, using (3.58) and reversing above arguments, it is clear that (3.54) or (3.55) also implies the associativity (1.52). $\blacksquare$


Let $V_{cl} = \oplus_{i=1}^N W_i^L \otimes W_i^R$. We define a map $T(V_{cl}) \boxtimes V_{op} \xrightarrow{\sigma_1} V_{op} \boxtimes T(V_{cl})$ by

$$\sigma_1 := \oplus_{i=1}^N (\mathcal{R}_+ \boxtimes \text{id}_{W_i^R}) \circ \mathcal{A} \circ (\text{id}_{W_i^L} \boxtimes \mathcal{R}_-) \circ \mathcal{A}^{-1}. \quad (3.61)$$

Proposition 3.10. *The commutativity I (recall Proposition 1.18) is equivalent to the following identity:*

$$\mu_{cl-op}(\text{id}_{T(V_{cl})} \boxtimes \mu_{op}) = \mu_{op}(\text{id}_{V_{op}} \boxtimes \mu_{cl-op}) \circ \mathcal{A}^{-1} \circ \sigma_1 \circ \mathcal{A}, \quad (3.62)$$

or the following graphic identities:



$$(3.63)$$

Proof. There is a morphism

$$\mu_{cl-op}^{P(z-\zeta)P(\zeta)} \circ (\text{id}_{T_{P(z-\zeta)}(V_{cl})} \boxtimes_{P(\zeta)} \mu_{op}^{P(r)}) \quad (3.64)$$

in $\text{Hom}(T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} (V_{op} \boxtimes_{P(r)} V_{op}))$ associated to (1.58). There is another morphism

$$\mu_{op}^{P(\tilde{r})} \circ (\text{id}_{V_{op}} \boxtimes_{P(\tilde{r})} \mu_{cl-op}^{P(z-\zeta)P(\zeta)}) \quad (3.65)$$

in $\text{Hom}(V_{op} \boxtimes (T(V_{cl}) \boxtimes V_{op}))$ associated to (1.59).

For z, ζ, r, r_1 in a proper subdomain of $r_1 > z > \zeta > r > 0$, using similar arguments as in Proposition 3.8, we obtain that the commutativity I, implies the following

commutative diagram:

$$\begin{array}{ccc}
T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} (V_{op} \boxtimes_{P(r)} V_{op}) & & (3.66) \\
\downarrow f_4 & \searrow & \\
T(V_{cl}) \boxtimes (V_{op} \boxtimes V_{op}) & & (3.64) \\
\downarrow \mathcal{A} & \searrow & \\
(T(V_{cl}) \boxtimes V_{op}) \boxtimes V_{op} & & V_{op} \\
\downarrow \sigma_1 & & \downarrow \text{id}_{V_{op}} \\
(V_{op} \boxtimes T(V_{cl})) \boxtimes V_{op} & & V_{op} \\
\downarrow \mathcal{A}^{-1} & \searrow & \\
V_{op} \boxtimes (T(V_{cl}) \boxtimes V_{op}) & & (3.65) \\
\downarrow g_4 & \nearrow & \\
V_{op} \boxtimes_{P(r_1)} (T_{P(z-\zeta)}(V_{cl}) \boxtimes_{P(\zeta)} V_{op}) & &
\end{array}$$

where

$$f_4 = (\mathcal{T}_{\gamma_2} \boxtimes \mathcal{T}_{\gamma_3}) \circ \mathcal{T}_{\gamma_1}, \quad (3.67)$$

$$g_4 = (\text{id}_{V_{op}} \boxtimes (\mathcal{T}_{\gamma_5} \boxtimes \mathcal{T}_{\gamma_6})) \circ \mathcal{T}_{\gamma_4}, \quad (3.68)$$

in which $\gamma_i, i = 1, 2, 3$ are paths in $\mathbb{R}_+$ from $\zeta, z - \zeta$ and r respectively to 1 and $\gamma_i, i = 4, 5, 6$ are path in $\mathbb{R}_+$ from 1 to $r_1, z - \zeta$ and ζ respectively. Above commutative diagram immediately implies that the subdiagram in the middle of (3.66) is commutative. This is nothing but the commutativity (3.62) or the first formula in (3.63). Moreover, it also easy to see that the two formula in (3.63) are actually equivalent.


Conversely, using commutative diagram (3.66) and reversing above arguments, it is clear that (3.63) implies the commutativity of rational $\tilde{\mathfrak{S}}^c$. $\blacksquare$

Commutativity II (recall Proposition 1.19) is not needed in Theorem 1.28 because it automatically follows from associativity II and skew symmetry of V_{cl} . It also has a very nice categorical formulation as given in the following proposition, which follows from (3.18) and (3.31) immediately.

Proposition 3.11. *For an open-closed field algebra over V , we have*

$$\mu_{cl-op} \circ \text{id}_{T(V_{cl})} \boxtimes \mu_{cl-op} = \mu_{cl-op} \circ \text{id}_{T(V_{cl})} \boxtimes \mu_{cl-op} \circ \mathcal{A}^{-1} \circ \sigma \circ \mathcal{A},$$

or equivalently



$$(3.69)$$

In summary, we have already completely reformulated the all the data and conditions in Theorem 1.28 in the language of tensor category as (3.23), (3.24), (3.30), (3.54) and (3.62) or equivalently as graphic identities (3.25), (3.31), (3.55) and (3.63).

We define a morphism $\iota_{cl-op} : T(V_{cl}) \rightarrow V_{op}$ as the composition of the following maps:

$$T(V_{cl}) \xrightarrow{r_{T(V_{cl})}^{-1}} T(V_{cl}) \boxtimes \mathbf{1}_{\mathcal{C}_V} \xrightarrow{\text{id} \boxtimes \iota_{V_{op}}} T(V_{cl}) \boxtimes V_{op} \xrightarrow{\mu_{cl-op}} V_{op}. \quad (3.70)$$

or equivalently as the following graphic formula:

$$\iota_{cl-op} = \begin{array}{c} | \\ \triangleup \\ | \end{array} := \begin{array}{c} | \\ \curvearrowright \triangle \\ | \end{array} \quad . \quad (3.71)$$

Lemma 3.12. ι_{cl-op} is an algebra morphism from $T(V_{cl})$ to V_{op} .

Proof. That ι_{cl-op} maps identity to identity is proved as follows:

$$\begin{array}{c} | \\ \triangleup \\ | \end{array} = \begin{array}{c} | \\ \curvearrowright \triangle \\ | \end{array} = \begin{array}{c} | \\ | \\ \circ \end{array} \quad . \quad (3.72)$$

The homomorphism property $\mu_{op} \circ (\iota_{cl-op} \boxtimes \iota_{cl-op}) = \iota_{cl-op} \circ \tilde{m}_{cl}$ is proven as follows:

$$\begin{array}{c} | \\ \triangleup \quad \triangleup \\ | \quad | \end{array} = \begin{array}{c} | \\ \curvearrowright \triangle \\ \curvearrowright \triangle \\ | \end{array} = \begin{array}{c} | \\ \triangle \\ \triangle \\ | \end{array} = \begin{array}{c} | \\ \triangleup \\ \triangle \\ | \end{array} \quad . \quad (3.73)$$

■

Definition 3.13. An open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebra, denoted as

$$((A_{op}, \mu_{op}, \iota_{op})|(A_{cl}, \mu_{cl}, \iota_{cl}), \iota_{cl-op})$$

or simply $(A_{op}|A_{cl}, \iota_{cl-op})$, consists of an associative algebra $(A_{op}, \mu_{op}, \iota_{op})$ in $\mathcal{C}_V$, a commutative associative algebra with a trivial twist $(A_{cl}, \mu_{cl}, \iota_{cl})$ in $\mathcal{C}_{V \otimes V}$ and an associative algebra homomorphism $\iota_{cl-op} : T(A_{cl}) \rightarrow A_{op}$, satisfying the following commutativity:

$$\mu_{op} \circ (\iota_{cl-op} \boxtimes \text{id}_{V_{op}}) = \mu_{op} \circ (\text{id}_{V_{op}} \boxtimes \iota_{cl-op}) \circ \sigma_1, \quad (3.74)$$

or equivalently,

$$\begin{array}{c} | \\ \triangleup \\ | \end{array} \quad \begin{array}{c} | \\ \triangleup \\ | \end{array} = \begin{array}{c} | \\ \triangleup \\ | \end{array} \quad \begin{array}{c} | \\ \triangleup \\ | \end{array} \quad . \quad (3.75)$$

Theorem 3.14. The notion of open-closed field algebra over V is equivalent to that of open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebra, in the sense that the categories of above two notions are isomorphic.

Proof. Given an open-closed field algebra over V , we have shown that it gives a triple $(V_{cl}, V_{op}, \mu_{cl-op})$, in which V_{cl} is a commutative associative algebra in $\mathcal{C}_{V \otimes V}$ with a trivial twist, and V_{op} is an algebra in $\mathcal{C}_V$, and μ_{cl-op} satisfies (3.25), (3.31), (3.55) and (3.63). Moreover, we have shown that ι_{cl-op} defined by (3.71) gives an morphism of associative algebra. Now we prove (3.75) as follows:

Hence $(V_{op}|V_{cl}, \iota_{cl-op})$ is an open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebra.

It is easy to check that it gives a functor from the category of open-closed field algebras over V to that of open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebras.

Conversely, given an open-closed $\mathcal{C}_V|\mathcal{C}_{V \otimes V}$ -algebra, $(V_{op}|V_{cl}, \iota_{cl-op})$, we define a morphism $\mu_{cl-op} \in \text{Hom}(T(V_{cl}) \boxtimes V_{op}, V_{op})$ as

$$\mu_{cl-op} = \text{[Diagram: two strands from the left, one straight, one curved, meeting at a triangle pointing right]} := \text{[Diagram: two strands from the bottom, one straight, one curved, meeting at a triangle pointing up]} . \quad (3.76)$$

Since ι_{cl-op} is an algebra homomorphism, it maps unit to unit (recall (3.72)). Thus the identity property of μ_{cl-op} holds. Then the identity property of $\mathbb{Y}_{cl-op}$ follows.

Furthermore, we have

which gives the associativity II (3.31). The associativity I (3.55) follows from

By using (3.2), we can prove the commutativity (3.63) as follows:

Notice that the other half of (3.63) is equivalent to the first half. Also notice that the commutativity (3.69) simply follows from (3.15) and the fact that ι_{cl-op} is an algebra homomorphism.

It is easy to check that this correspondence gives a functor from the category of open-closed $\mathcal{C}_V|\mathcal{C}_{V\otimes V}$ -algebras to that of open-closed field algebras over V .

That the two relevant categories are isomorphic follows from (3.71) and (3.76) easily. ■

It is very easy to construct open-closed $\mathcal{C}_V|\mathcal{C}_{V\otimes V}$ -algebras. For example, let A be an associative algebra in $\mathcal{C}_{V\otimes V}$ and $C_l(A)$ the left center of A [O]. Let A_0 be any subalgebra of $C_l(A)$ and $\iota : A_0 \hookrightarrow A$ the natural embedding. Then it is clear that $(T(A)|_{A_0}, T(\iota))$ gives an open-closed $\mathcal{C}_V|\mathcal{C}_{V\otimes V}$ -algebra, which further gives an open-closed field algebra over V and a smooth $\tilde{\mathfrak{S}}^c$ -algebra over V . We will not pursue the construction further in this work. Instead, we leave it to [K3], in which we will construct explicit examples that are most relevant in physics. Moreover, a general theory of open-closed $\mathcal{C}_V|\mathcal{C}_{V\otimes V}$ -algebras and its variants ([K3]) will be developed and its connection to the works [FFFS][FS3][FRS1]-[FRS4][FjFRS] will be explained in future publications.

References

- [A] V. Alessandrini, A general approach to Dual Multiloop Diagrams, *Il Nuovo Cimento*, Vol 2A. N. 2, (1971).
- [B] R. E. Borcherds, Quantum Vertex algebras, *Taniguchi Conference on Mathematics Nara'98*, 51–74, *Adv. Stud. Pure Math.*, 31, *Math. Soc. Japan.*, Tokyo, 2001.
- [BKa] B. Bakalov and V. Kac, Field algebras *Int. Math. Res. Not.* **2003**, no. 3, 123-159.
- [BKl] B. Bakalov and A. Kirillov Jr., *Lectures on Tensor Categories and Modular Functors*, University Lecture Series, Vol. 21, Amer. Math. Soc., Providence, 2001.
- [C1] J. L. Cardy, Conformal invariance and surface critical behavior, *Nucl. Phys.* **B240** (1984), 514-532.
- [C2] J. L. Cardy, Effect of boundary conditions on the operator content of two-dimensional conformally invariant theories, *Nucl. Phys.* **B275** (1986), 200–218.
- [C3] J. L. Cardy, Operator content of two-dimensional conformal invariant theories, *Nucl. Phys.* **B270** (1986) 186-204.
- [C4] J. L. Cardy, Boundary conditions, fusion rules and the Verlinde formula, *Nucl. Phys.* **B324** (1989), 581–596.
- [CL] J. Cardy and D. C. Lewellen, Bulk and boundary operators in conformal field theory, *Phys. Lett.* **B259** (1991) 274.
- [EK] P. Etingof and D. Kazhdan, Quantization of Lie bialgebras. V. Quantum vertex operator algebras, *Selecta Math.* (N.S.) **6** (2000), 105-130.

- [FFFS] G. Felder, J. Fröhlich, J. Fuchs and C. Schweigert, Correlation functions and boundary conditions in rational conformal field theory and three-dimensional topology, *Compositio Math.* **131** (2002), 189–237.
- [FjFRS] J. Fjelstad, J. Fuchs, I. Runkel and C. Schweigert, TFT construction of RCFT correlators V: Proof of modular invariance and factorisation, hep-th/0503194.
- [FHL] I. B. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Memoirs Amer. Math. Soc.* **104**, 1993.
- [FR] E. Frenkel and N. Reshetikhin, Towards deformed chiral algebras, in Proc. Quantum Group Symp. at XXIth Intern. Coll. on Group Theoretical Method in Physics, Goslar 1996.
- [FRS1] J. Fuchs, I. Runkel and C. Schweigert, Conformal correlation functions, Frobenius algebras and triangulations, *Nuc Phys.* **B624** (2002), 452–468.
- [FRS2] J. Fuchs, I. Runkel and C. Schweigert, TFT construction of RCFT correlators I: Partition functions, *Nucl. Phys.* **B678** (2004) 511.
- [FRS3] J. Fuchs, I. Runkel and C. Schweigert, TFT construction of RCFT correlators III: Simple currents, *Nucl. Phys.* **B694** (2004) 277.
- [FRS4] J. Fuchs, I. Runkel and C. Schweigert, TFT construction of RCFT correlators IV: Structure constants and correlation functions, preprint hep-th/0412290, to appear in Nucl. Phys. B.
- [FS1] J. Fuchs and C. Schweigert, Symmetry breaking boundaries I. General theory, *Nucl. Phys.* **B558** (1999) 419–483.
- [FS2] J. Fuchs and C. Schweigert, Symmetry breaking boundaries II. More structures; Examples *Nucl. Phys.* **B568** (2000) 543–593.
- [FS3] J. Fuchs and C. Schweigert, Category theory for conformal boundary conditions, *Fields Institute Commun.* 39 (2003) 25.
- [H1] Y.-Z. Huang, *On the geometric interpretation of vertex operator algebras*, Ph.D thesis, Rutgers University, 1990.
- [H2] Y.-Z. Huang, Geometric interpretation of vertex operator algebras, *Proc. Natl. Acad. Sci. USA* **88** (1991), 9964–9968.
- [H3] Y.-Z. Huang, A theory of tensor products for module categories for a vertex operator algebra, IV, *J. Pure Appl. Alg.* **100** (1995), 173–216.
- [H4] Y.-Z. Huang, *Two-dimensional conformal geometry and vertex operator algebras*, Progress in Mathematics, Vol. 148, Birkhäuser, Boston, 1997.
- [H5] Y.-Z. Huang, Intertwining operator algebras, genus-zero modular functors and genus-zero conformal field theories, in *Operads: Proceedings of Renaissance Conferences*, ed. J.-L. Loday, J. Stasheff, and A. A. Voronov, Contemporary Math., Vol. 202, Amer. Math. Soc., Providence, 1997, 335–355.

- [H6] Y.-Z. Huang, Genus-zero modular functors and intertwining operator algebras, *Internat. J. Math.* **9** (1998), 845-863.
- [H7] Y.-Z. Huang, Riemann surfaces with boundaries and the theory of vertex operator algebras, *Fields Institute Commun.* **39** (2003) 109.
- [H8] Y.-Z. Huang, Differential equations and intertwining operators, *Comm. Contemp. Math.* **7** (2005), 375-400.
- [H9] Y.-Z. Huang, Vertex operator algebras and the Verlinde conjecture, *Comm. Contemp. Math.*, to appear; math.QA/0406291.
- [HKo1] Y.-Z. Huang and L. Kong, Open-string vertex algebra, category and operad, *Comm. Math. Phys.*, **250** (2004), 433-471.
- [HKo2] Y.-Z. Huang and L. Kong, Full field algebra, *Comm. Math. Phys.*, **272** (2007) 345-396.
- [HKo3] Y.-Z. Huang and L. Kong, Modular invariance for full field algebras, math.QA/0609570.
- [HKr] P. Hu and I. Kriz, Closed and open conformal field theories and their anomalies, *Comm. Math. Phys.* **254** (2005), no. 1, 221-253.
- [HL1] Y.-Z. Huang and J. Lepowsky, Vertex operator algebras and operads, *The Gelfand Mathematical Seminars, 1990-1992*, ed. L. Corwin, I. Gelfand and J. Lepowsky, Birkhäuser, Boston, 1993, 145-161.
- [HL2] Y.-Z. Huang and J. Lepowsky, Tensor products of modules for a vertex operator algebra and vertex tensor categories, in: *Lie Theory and Geometry, in honor of Bertram Kostant*, ed. R. Brylinski, J.-L. Brylinski, V. Guillemin, V. Kac, Birkhäuser, Boston, 1994, 349-383.
- [HL3] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, I, *Selecta Math. (N.S.)* **1** (1995), 699-756.
- [HL4] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, II, *Selecta Math. (N.S.)* **1** (1995), 757-786.
- [HL5] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, III, *J. Pure Appl. Alg.* **100** (1995), 141-171.
- [K1] L. Kong, Nondegenerate intertwining operator algebras, unpublished.
- [K2] L. Kong, Full field algebras, operads and tensor categories, *Adv. Math* **213** (2007) 271-340.
- [K3] L. Kong, Cardy condition for open-closed field algebras, math.QA/0612255.
- [Kt] M. Kontsevich, Operads and motives in deformation quantization, *Letters in Mathematical Physics* **48**: 35-72, (1999).
- [L1] H.S. Li, Axiomatic G_1 -vertex algebras, *Commun. Contemp. Math.* **5** (2003), 281-327.

- [L2] H.S. Li, Nonlocal vertex algebras generated by formal vertex operators, *Selecta Math. (N.S.)* **11** (2005), no. 3-4, 349–397.
- [L3] H.S. Li, Constructing quantum vertex algebras, *Internat. J. Math.* **17** (2006), no. 4, 441–476.
- [MSS] M. Markl, S. Shnider and J. D. Stasheff, *Operads in Algebra, Topology and Physics*, volume 96 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, Rhode Island, 2002.
- [O] V. Ostrik, Module categories, weak Hopf algebras and modular invariants, *Transform. Groups* **8** (2003), 177–206.
- [S1] G. Segal, The definition of conformal field theory, in: *Differential geometrical methods in theoretical physics (Como, 1987)*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 250, Kluwer Acad. Publ., Dordrecht, 1988, 165–171.
- [S2] G. Segal, The definition of conformal field theory, preprint, 1988; also in: *Topology, geometry and quantum field theory*, ed. U. Tillmann, London Math. Soc. Lect. Note Ser., Vol. 308. Cambridge University Press, Cambridge, 2004, 421–577.
- [V] A. A. Voronov, The Swiss-cheese operad, in: *Homotopy invariant algebraic structures, in honor of J. Michael Boardman, Proc. of the AMS Special Session on Homotopy Theory, Baltimore, 1998*, ed. J.-P. Meyer, J. Morava and W. S. Wilson, Contemporary Math., Vol. 239, Amer. Math. Soc., Providence, RI, 1999, 365–373.

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, D-04103, LEIPZIG, GERMANY

and

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, LE BOIS-MARIE, 35, ROUTE DE CHARTRES, F-91440 BURES-SUR-YVETTE, FRANCE

and

MAX-PLANCK-INSTITUT FÜR MATHEMATIK VIVATSGASSE 7, D-53111 BONN, GERMANY (current address)

E-mail address: kong@mpim-bonn.mpg.de